

Accepted Manuscript

Assembly and disassembly of *Aspergillus fumigatus* conidial rodlets

Isabel Valsecchi, Jennifer I. Lai, Emmanuel Stephen-Victor, Ariane Pillé, Audrey Beaussart, Victor Lo, Chi L.L. Pham, Vishukumar Aimanianda, Ann H. Kwan, Magalie Duchateau, Quentin Gai Gianetto, Mariette Matondo, Melanie Lehoux, Donald C. Sheppard, Yves F. Dufrene, Jagadeesh Bayry, J. Iñaki Guijarro, Margaret Sunde, Jean-Paul Latgé


PII: S2468-2330(18)30022-7
DOI: <https://doi.org/10.1016/j.tcsw.2019.100023>
Article Number: 100023
Reference: TCSW 100023

To appear in: *The Cell Surface*

Received Date: 12 June 2018
Revised Date: 22 February 2019
Accepted Date: 23 February 2019

Please cite this article as: I. Valsecchi, J.I. Lai, E. Stephen-Victor, A. Pillé, A. Beaussart, V. Lo, C.L.L. Pham, V. Aimanianda, A.H. Kwan, M. Duchateau, Q.G. Gianetto, M. Matondo, M. Lehoux, D.C. Sheppard, Y.F. Dufrene, J. Bayry, J.I. Guijarro, M. Sunde, J-P. Latgé, Assembly and disassembly of *Aspergillus fumigatus* conidial rodlets, *The Cell Surface* (2019), doi: <https://doi.org/10.1016/j.tcsw.2019.100023>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Assembly and disassembly of *Aspergillus fumigatus* conidial rodlets

Isabel Valsecchi^{a,d,*}, Jennifer I. Lai^{b,*}, Emmanuel Stephen-Victor^{c*}, Ariane Pillé^{d*}, Audrey Beaussart^{e,f,1*}, Victor Lo^b, Chi L.L. Pham^b, Vishukumar Aimanianda^a, Ann H. Kwan^g, Magalie Duchateau^h, Quentin Gai Gianetto^{h,i}, Mariette Matondo^h, Melanie Lehoux^j, Donald C. Sheppard^j, Yves F. Dufrene^{e,1#}, Jagadeesh Bayry^{c#}, J. Iñaki Guijarro^{d,1#}, Margaret Sunde^{b#}, Jean-Paul Latgé^{a#}

^a Unité des Aspergillus, Institut Pasteur, Paris, France

^b School of Medical Sciences and Sydney Nano, The University of Sydney, NSW 2006, Australia

^c Institut National de la Santé et de la Recherche Médicale; Centre de Recherche des Cordeliers; Sorbonne Université, Paris, France

^d Biological NMR Technological Platform, Institut Pasteur, CNRS UMR 3528, Paris, France

^e Institute of Life Sciences, Université Catholique de Louvain, Croix du Sud, 4-5, bte L7.07.06, B-1348 Louvain-la-Neuve, Belgium

^f Walloon Excellence in Life Sciences and Biotechnology, Belgium

^g School of Life and Environmental Sciences and Sydney Nano, The University of Sydney, NSW 2006, Australia

^h Pasteur Proteomics Platform, Mass Spectrometry for Biology Unit, Institut Pasteur, CNRS USR 2000, Paris, France

ⁱ Bioinformatics and Biostatistics Hub, C3BI, CNRS USR 3756, Institut Pasteur, Paris, France

^j Departments of Medicine, Microbiology and Immunology, McGill University, Montréal, QC, Canada.

¹ Present address: Université de Lorraine, CNRS, LIEC, 54000 Nancy, France.

* Co-first authors, have contributed equally to the study

Joint corresponding authors

Assembly and disassembly of *Aspergillus fumigatus* conidial rodlets

Abstract

The rodlet structure present on the *Aspergillus fumigatus* conidial surface hides conidia from immune recognition. In spite of the essential biological role of the rodlets, the molecular basis for their self-assembly and disaggregation is not known. Analysis of the soluble forms of conidia-extracted and recombinant RodA by NMR spectroscopy has indicated the importance of disulfide bonds and identified two dynamic regions as likely candidates for conformational change and intermolecular interactions during conversion of RodA into the amyloid rodlet structure. Point mutations introduced into the *RODA* sequence confirmed that (1) mutation of a single cysteine was sufficient to block rodlet formation on the conidial surface and (2) both presumed amyloidogenic regions were needed for proper rodlet assembly. Mutations in the two putative amyloidogenic regions retarded and disturbed, but did not completely inhibit, the formation of the rodlets *in vitro* and on the conidial surface. In contrast to the parental conidia, mutant conidia lacking disulfide bridges within RodA or expressing RodA carrying the double (I115S/I146G) mutation activated dendritic cells with the subsequent secretion of proinflammatory cytokines. The immune reactivity of the RodA mutant conidia was not due to a modification in the RodA structure, but to the exposure of different pathogen-associated molecular patterns on the surface as a result of the modification of the rodlet surface layer. The disorganization and loss of the rodlet layer which occur during the early stages of germination, and lead to an anti-*fumigatus* host immune response, is initiated by the cell wall bound aspartic proteases.

Keywords

Aspergillus fumigatus; hydrophobins; rodlets; RodA; amyloids.

Introduction

Aerial conidia of Ascomycetes and Basidiomycetes are coated with a layer of hydrophobin proteins organized in fibrillar structures known as rodlets. The rodlet layer is amphipathic, and the outward facing hydrophobic surface renders the conidial surface resistant to wetting, thus facilitating effective dispersal of conidia in the air (Gebbinck et al., 2005; Sunde et al., 2008). Fungal hydrophobins are small secreted proteins with eight cysteine residues ordered in a distinct way and with characteristic hydrophathy patterns, which define two well-characterized classes I and II and a third intermediate class named III (Gandier et al., 2017; Kershaw and Talbot, 1998; Littlejohn et al., 2012; Pedersen et al., 2011). Class I hydrophobin assemblies are generally water- and detergent-insoluble, have an amyloid structure and form the patterned rodlet layer observed in conidia. Class II hydrophobin sequences are generally shorter than class I hydrophobin sequences and are soluble in heated aqueous ethanol mixtures and SDS. The intermediate class III has the same cysteine pattern but lacks a consistent hydrophathy profile characteristic of either class I or class II hydrophobins (Jensen et al., 2010; Littlejohn et al., 2012).

In addition to the role of the rodlet layer in fungal life during conidiogenesis and conidial dispersal in the air, it has been shown that rodlets have a role in fungal–host interactions and favor the pathogenic behaviour of fungal pathogens in mammals, plant and insect hosts (Aimanianda et al., 2009; Talbot et al., 1996; Zhang et al., 2011) or play a positive role during plant symbiosis (Whiteford and Spanu, 2002). Moreover, due to their specific physicochemical properties, hydrophobins have potential for numerous biotechnological applications (Hektor and Scholtmeijer, 2005; Piscitelli et al., 2017; Zhao et al., 2009). Their coating properties could be used to functionalize metals and plastic or carbon nanotubes in order to prevent the formation of bacterial biofilms or to immobilize enzymes on surface (Tanaka et al., 2017) or to stabilize pharmaceutical emulsions and facilitate drug delivery (Artini et al., 2017).

Our current study is focused on the RodA hydrophobin of *A. fumigatus*, the major airborne opportunistic fungal pathogen of humans. This hydrophobin forms rodlets on the surface of *A. fumigatus* conidia (Paris et al., 2003; Thau et al., 1994). This surface rodlet coat masks the conidium from host detection and its removal from the conidial surface initiates the antifungal host immune response (Aimanianda et al., 2009). Despite their prominent morphological and biological role, the molecular basis for rodlet self-assembly and disaggregation in this fungal species has not yet been understood. By analysing the structure of the soluble form of RodA and by creating point mutations in the *RODA* gene, we report here on the mechanism that drives RodA to assemble into highly ordered structures anchored to the conidial surface. Moreover, we have also investigated the biochemical events responsible for the lysis of the rodlet layer, a key step in the recognition and mounting of an immune response to this airborne pathogen by host cells.

Results

Solution structure and dynamics of RodA

The structure of RodA was studied by NMR spectroscopy. Comparison of the ^1H - ^{15}N HSQC spectra of recombinant RodA (rRodA) and the HPLC purified hydrofluoric acid-extracted RodA from conidia indicated that both proteins have the same structure, the same cysteine redox state and potential disulfide pairing (Supplementary Material Fig. S1). Additionally, the native protein did not contain any post-translational modification such as an O- or N-glycosylation. This was confirmed by mass spectrometry (MS) of proteolysis fragments of the

full-length conidia-extracted protein before or after chemical treatment with the N- and O-deglycosylating agent trifluoromethanesulfonic acid (data not shown). We thus performed further work using the recombinant protein expressed in *E. coli* that could be purified in greater amounts than the native conidial RodA. Analysis of the cysteine redox state as proposed by Sharma Rajarathnam (Sharma and Rajarathnam, 2000) indicated that the Cys residues of rRodA were oxidized and hence, involved in disulfide bridges. Distance (nOe) data and topology of rRodA allowed us to unambiguously establish the disulfide pairing of rRodA, which corresponded to the one observed in all other studied hydrophobins (C1-C6: 56-133; C2-C5: 64-127; C3-C4: 65-105; C7-C8: 134-152) (Fig.1). Cysteines are numbered according to their order of appearance in the sequence from C1 to C8 or by their sequence number throughout the manuscript.

The structure of RodA (Fig. 1 and Supplementary Table S1) is organized around the four-disulfide bonds. It consists of a central β -barrel composed of two curved antiparallel β -sheets (sheet S1: strands 60-65, 104-107, 130-135 and 150-154; sheet S2: strands 65-70, 99-104 and 150-154), two relatively long α -helices (helix H1: 49-56; H2: 79-90), one short α -helix (H4: 123-126) and two short 3_{10} - (H3: 96-98; H5: 155-157) helices. Two long and kinked β -strands (60-70 and 99-107) participate in both β -sheets of the β -barrel. Two S-S bonds link together strands of the β -barrel and the other two connect the external face of the β -barrel with the regions of helices H1 and H4. The long α -helix H2 in the C3-C4 region packs against the β -barrel. The structures showed good convergence within the regions with secondary structure, but were more variable for residues 19-39, 109-121 and 137-149 located in the N-terminal, C4-C5 (106-132) and C7-C8 (135-151) inter-cysteine regions, respectively (Fig. S2).

The internal motions of rRodA on the nanosecond-picosecond time scale were analyzed using the backbone amide ^1H - ^{15}N nOes (Fig. 1B and Supplementary Fig. S3). Low nOe values are indicative of high amplitude internal motions while high nOe values reflect rigidity of the backbone at the fast timescale (ns-ps). The nOe data showed that the secondary structure elements of RodA in the vicinity of disulfide bridges, namely the central β -barrel and α -helices H1 and H4 that are tethered to the β -barrel by S-S bonds are rigid with nOe values higher than 0.71. Helix H2, located in the region between cysteine residues C3 and C4, showed somewhat lower ^1H - ^{15}N nOe values (0.56 to 0.85) suggestive of low amplitude motions. In contrast, the N-terminal region (19-39) showed negative or very low nOes that indicated that this region was disordered. Residues 109-121 (C4-C5 region), 137-149 (C7-C8 region) and to a lesser extent the flanking regions (74-79 and 90-96) of α -helices H2 and H3 in the C3-C4 region also display low nOe values.

The charge distribution and hydrophobicity of the surface of hydrophobins is important for the recruitment and self-assembly of hydrophobins at a hydrophobic/hydrophilic interface such as the air/water interface (Fig. 1C). The surface of RodA contains a hydrophobic region between C3 and C4 and two highly hydrophobic patches (114-120 and 136-148) in the inter-cysteine regions C4-C5 and C7-C8. A relatively hydrophobic belt joins the hydrophobic and flexible C4-C5 and C7-C8 loops. The belt is lined by charged residues with two pairs of charged residues (E54-K50; K107-D109) that establish salt bridges and compensate their charges. An opposite face of the molecule shows many charged residues with two clusters of negative residues (D73, D76, D78 and E79 in the C3-C4 region and D45 and D46 close to the disordered N-terminal tail) and isolated positive residues (K67, K128 and K55 and K87). Hence, in solution, monomeric RodA shows an amphiphilic character with a hydrophobic face with hydrophobic residues and compensated charges and a more hydrophilic surface with clusters of net charges on the opposite face.

Molecular identification of the amyloidogenic regions within RodA protein and the amino acids essential for the formation of the rodlets *in vitro*

Analysis of the RodA sequence by the consensus method Amylpred2 (Tsolis et al., 2013) indicated that stretches of residues (112-117 and 143-147) within the C4-C5 and C7-C8 inter-cysteine regions are predicted as amyloidogenic and could potentially participate in the cross β -sheets at the core of amyloid fibres. To investigate the role of these sequences in the context of the intact protein and interface-driven self-assembly, a number of mutations were introduced into rRodA. Individual or multiple glycine and serine substitutions of isoleucine and leucine residues were used to reduce side chain hydrophobicity and size and to introduce flexibility or polarity at sites within the protein structure. The kinetics of self-assembly of rRodA from the monomeric form into the amyloid rodlet form were monitored by changes in fluorescence of the amyloid binding dye Thioflavin T (ThT) during *in vitro* self-assembly assays (Fig. 2).

Only mutations to the rRodA sequence in the C4-C5 loop or in the C7-C8 loop resulted in a delay in self-assembly of rRodA mutants, relative to wild-type (WT) rRodA. Single-residue mutations introduced a significant lag phase of 10–20 min into the self-assembly profile, whereas no lag phase was detected for WT rRodA under these conditions (Fig. 2). The effect of double mutations in the C4–C5 and C7–C8 loops was additive since a rRodA carrying the double slowing mutations I115G/I146G displayed a longer lag phase than rRodA I115G and rRodA I146G (Fig.2). Mutations introduced in other regions of rRodA outside of the C4–C5 and C7–C8 loops did not have an effect on RodA self-assembly (data not shown). Strikingly, a chimeric protein rRodA C7C8NC2, which has the central sequence of the rRodA C7–C8 loop replaced with the C7–C8 loop from the non-amyloidogenic class II hydrophobin NC2, was able to assemble into ThT-positive structures (Fig. 2). A similar chimera in which the NC2 region was inserted into the hydrophobin EAS did not produce ThT-positive rodlets. These data show that in contrast to the hydrophobin EAS where only the C7–C8 loop is required for rodlet formation (Macindoe et al., 2012), RodA rodlet assembly involves the C4–C5 and the C7–C8 loops.

As revealed by atomic force microscopy (AFM) images in Fig. 3A, WT rRodA rodlets have comparable dimensions to those reported for other class I hydrophobins: width of 10.1 ± 1.8 nm and height of 2.0 ± 0.5 nm (Table S2). Introduction of the single mutations between cysteines 4 and 5 (Fig. 3 B,C) and cysteines seven and eight (Fig. 3 D,E) that slowed assembly kinetics did not grossly alter the morphology of the final assembled rodlet structure on a hydrophobic graphite surface (HOPG), as probed by AFM. In contrast, the rRodA protein carrying the double mutation and the rRodA C7C8NC2 chimera did not spontaneously form rodlets when dried onto HOPG at room temperature. However, when rRodA I115G/I146G protein was incubated at 50°C for 2 hours on the graphite surface, it formed a film composed of very long rodlets, as did WT rRodA assembled at this temperature (Fig. 3F, G). The rRodA C7C8NC2 chimera also formed rodlets only when prepared at 50°C. However, the surface details and lateral packing of the rodlets formed by the chimera were less distinct than observed with WT rRodA rodlets (Fig. 3H).

The results obtained with the different point mutated rRodA proteins suggest that while a single amyloidogenic region can support the incorporation of monomers into a fibrillar structure albeit with slower kinetics, the absence of a second interaction within the rodlets causes a morphological disturbance, with consequences for the organization and lateral interactions between rodlets within the film. The self-assembling nature of

the sequences between the 4-5th and 7-8th cysteine residues was confirmed by analysis of peptides containing these sequences. Transmission electron microscopy showed that both the peptide PIIGIQDL (Fig. 4A) that originates from the C4-C5 region and SLIGL (Fig. 4B) found in the C7-C8 region can form fibrillar structures. In PIIGIQDL, fibrillar aggregation occurs at a concentration of 200 $\mu\text{g/ml}$, with many smaller fibrils bundling together to form a single fibril. The SLIGL peptide formed thicker single fibrils with a crystalline nature at 200 $\mu\text{g/ml}$.

Assembly of the rodlet structure on the conidium

In order to determine the effect of RodA mutations on rodlet assembly on conidia, different mutations were also introduced into the *RODA* gene (Figs. S4, S5). The effect of these mutations on the rodlet structure on the conidial surface was investigated with AFM (Figs. 5-7). Incubation of parental strain conidia with hydrofluoric acid (HF) or formic acid (FA) to isolate native RodA always results in the isolation of two protein bands (Aimanianda et al., 2009). MS, N-terminal sequencing and NMR data led to the identification of both molecular species. The higher M_r band corresponds to the full-length protein without the signal peptide with an N-terminal pyroglutamic acid (PCA) arising from a spontaneous deamidation of the N-terminal Gln residue at position 21 (PCA21-L159). The lower M_r species corresponds to the F41-L159 peptide, missing 40 amino acids from the N-terminus. The mutant with a RodA protein corresponding to the band with lower M_r seen in conidial extracts did not show any alteration in rodlet formation on the surface of the conidium (Fig. S4). This result indicates that the N-terminal region, which is unstructured in the monomer, does not play a role in the assembly of the rodlet layer.

The necessity for intact disulfide bridges within RodA for correct rodlet assembly in *A. fumigatus* was evaluated by the construction of mutants with the *RODA* gene mutated in one cysteine (C127S) or in four of them (C64S/C65S/C133S/C134S), disrupting one or all the four disulfide bridges. None of these mutants produced rodlets on the conidial surface as shown by AFM (Fig. 5). Moreover, negative Western blots of the formic acid extracts of these cysteine mutants with an anti-RodA antibody demonstrated that RodA was not secreted to the conidial surface (Fig. S6).

A. fumigatus mutants harboring the same point mutations as the ones undertaken with the recombinant protein (I114G, I115S, L145S, I146G and I115S/I146G) were generated. Like in the parental strain, RodA could be detected on SDS-PAGE as doublets after HF or FA extraction from all these RodA loop region mutants (Fig. S7). The surface organization of mutants resulting from the point mutations I114G, I115S, L145S, I146G and I115S/I146G was investigated by AFM. The hydrophobicity of the rodlet structure of the RodA I114G, RodA I115S and RodA L145S mutants was not affected, as seen by the adhesion force histograms obtained using chemical force microscopy with hydrophobic AFM tips (Dague et al., 2008b, 2007), data not shown]. In agreement with the ThT data obtained with mutated rRodA in *in vitro* assays (Fig. 2), the single-point mutation I146G and the double mutation I115S/I146G, which both led to a longer lag phase, affected the assembly of the rodlet layer on the surface of the conidium (Fig. 6). The appearance of the rodlets on the surface of the conidium was dependent on the conidial age. At 3-days of growth, conidia of these two mutants did not show any rodlets on their surface unlike parental conidia; however, the number of rodlets increased with the age of the conidia. The maximal rodlet formation was seen after 30-days of culture without further evolution of the rodlet morphology afterwards. Interestingly, two modifications of the rodlet structure were seen on the mutant conidia: (1) the length of the rodlets significantly increased from 147 nm observed for the parental to 270

and 234 nm for the I146G and I115S/I146G mutants, respectively (Fig. S8); (2) while parental rodlets coated the entire conidium, the coverage of the conidium surface by the mutated rodlets was incomplete, with a coverage ranging from 20 to 50% of the conidial surface for the I115S/I146G mutant (Fig. 7).

In addition, the surfaces of the conidia of the I146G and I115S/I146G mutants were covered at least partially by an amorphous material. The MS analysis of the extract obtained after incubation of the conidia in formic acid for 2 hours indicated that over 1200 proteins were quantified in the I115S/I146G mutant conidia and this number greatly exceeded the number of proteins detected with its parental strain, demonstrating the presence of a much higher concentration of proteins in the mutant cell wall (Data are available via ProteomeXchange with the identifier PXD008503). In spite of differences seen with AFM, a comparable number of proteins were extracted from 10-days or 30-day old I115S/I146G mutant conidia (1399 and 1278 respectively). The identification of the mutated and wild type RodA protein in the extracts from the mutant and parental strains validated the MS experiment. Analysis of proteins with a signal peptide, which are specifically targeted to be secreted to the cell wall, showed that 56 and 68 proteins were present in the cell wall of the 10 and 30-days old conidia, respectively (Table 1). Among the 31 proteins extracted from both 10- and 30-days mutant conidia, some of them such as disulfide isomerases, carboxypeptidases or phiA (Melin et al., 2003; Monod et al., 2002) could be specifically involved in the maturation of RodA. Others such as glucanases, chitinases, transglycosidases or GPI proteins are known to be involved in cell wall modifications (Latgé et al., 2017). Among these proteins, several (Sun1, Crf1, MP1, ChiB1, Aspf4, DPPV) are also known as antigens/allergens in agreement with their cell wall surface localization (Beauvais et al., 1997; Gastebois et al., 2013; Latgé, 1999). The coating of the outer conidial layer by an amorphous material was confirmed by the positive labelling of the conidia of these two mutants with the wheat germ agglutinin and concanavalin A lectins, in contrast with the parental strain (Fig. S9). The hydrophilicity of the I115S/I146G double mutant conidia seen by AFM was due to an incomplete coverage of the conidial surface by the rodlets and to the presence of amorphous (glyco)proteins on the surface of the mutant cell wall. Accordingly, the I115S/I146G mutant conidial colonies appeared black and hydrophilic like the $\Delta rodA$ mutant, as did (but to a lesser extent) the conidia from the single I146G mutant (Fig. S10). Complementation of the mutated *RODA* gene with the wild type gene led to the production of normal hydrophobic conidia displaying the characteristic green color of the parental strain instead of the black color of the mutant conidia (Fig. S10).

In conclusion, the results obtained with the conidial mutants were very reminiscent of the ones obtained *in vitro* with mutated recombinant RodA analyzed by ThT fluorescence assays in solution and AFM on a solid support. The rodlets of the mutants were organized differently when amino acids from both amyloidogenic regions were simultaneously mutated and/or when the protein carried a mutation that significantly extended the lag phase for rodlet formation. Although rodlets were still formed, their structural maturation required a longer time and was incomplete, leaving the conidia hydrophilic and with amorphous materials covering their surfaces.

Impact of point mutations on the immune response towards hydrophobins

We have reported earlier that the surface rodlet layer masks *A. fumigatus* conidial recognition by the host immune cells, whereas the conidial morphotypes devoid of a rodlet layer (i.e., $\Delta rodA$ mutant conidia or conidia after HF treatment) activate innate as well as adaptive immune responses (Aimanianda et al., 2009). These results prompted us to investigate if the

disorganization of the rodlet structure on the conidial surface as a consequence of point mutations in the *RODA* gene would lead to the activation of immune cells by mutant conidia. Since dendritic cells (DC) are major players in deciding the fate of the immune response, the effect of different *RODA*-mutant conidia was investigated on human DCs.

A. fumigatus conidia bearing mutations in the cysteine residue(s) [both single C127S or quadruple (C64S/C65S/C133S/C134S)] or the double I115S/I146G mutant conidia were indeed immunogenic. A significantly higher expression of CD83, a terminal maturation marker of DC, and of the co-stimulatory molecule CD86 was seen when DCs were exposed to cysteine (both single and quadruple) or I115S/I146G mutant conidia (Fig. 8A). Other DC maturation markers such as CD40, CD86 and HLA-DR were also enhanced by these mutant conidia (data not shown). In contrast, conidia with single mutations in the C4-C5 or C7-C8 loop regions (i.e., I114G, I115S, L145S, I146G point-mutant conidia) did not induce a DC response (Fig. 8B). Moreover, conidia with a cysteine mutation (both single or quadruple) or the double I115S/I146G mutation induced significantly higher quantities of the DC cytokines TNF α and IL6 (Figs. 8C,D). Untreated immature DCs produced only minimal quantities of cytokines like the parental or the single point mutated I114G, I115S or L145S conidia, while the I146G mutant conidia induced moderate amounts of these cytokines.

CD4⁺ T cell polarization mediated by *A. fumigatus* *RODA* point-mutant conidia matured DCs was also investigated. Quantification of secretion of T cell cytokines (IFN- γ , IL-4, IL-17A) or intracellular staining of the transcription factor FoxP3 revealed that cysteine mutant conidia [i.e., both C127S and C64S/C65S/C133S/C134S) mutant conidia] or harboring the double mutation (I115S/I146G) promoted distinct T cell responses. Single and quadruple cysteine mutant conidia-matured-DCs polarized predominantly CD4⁺CD25⁺CD127⁻ FoxP3⁺ Treg and IFN- γ (Th1) responses (Fig. 8E), whereas Th2 (IL-4) and Th17 (IL-17A) cytokines remained at basal levels (Fig. 8E). On the contrary, I115S/I146G double mutant conidia matured-DCs promoted only Th1 responses, while other T cell subsets, including Tregs, were not altered (Fig. 8F). Hence, *A. fumigatus* conidia that lack disulfide bridges in RodA or with the double mutation (I115S/I146G) in the amyloidogenic regions are both immunogenic but promoted distinct T cell responses.

Earlier, we had shown that RodA extracted from the conidial surface did not stimulate an immune response of the host (Aimanianda et al., 2009). Recombinant RodA, whether folded or unfolded, with non-reduced or reduced and blocked cysteines with N-ethyl maleimide (NEM) or iodoacetamide (IA), did not modify either the DC phenotype or the production of cytokines (Fig. 9A,B). Similar results were obtained for conidia-extracted RodA harboring single or double mutations in their C4-C5 or C7-C8 regions or lacking the forty N-terminal amino acids (Fig. 9C,D). These observations clearly indicated that the DC activation and maturation induced by the double I115S/I146G mutant conidia was only due to the emergence of different pathogen-associated molecular patterns on the surface of conidia, concomitant with the delayed formation of the rodlets and hence with the disorganization of the outer cell wall layer.

Disassembly of the rodlet structure during germination

In nature, the activation of the immune response is associated with the loss of the rodlet layer, which occurs during conidial germination (Aimanianda et al., 2009). However, the mechanism of disintegration of the rodlet layer during germination has not been previously investigated. Disassembly of the rodlet structure is a progressive event during conidial swelling. Complete removal of the rodlet structure from the conidial surface occurs in about

5-6 h from the commencement of the germination process (data not shown). When recombinant or native RodA proteins were incubated with the ethanol precipitated protein mixture from the supernatant of germinating conidia, RodA was degraded (Fig. S11). RodA degradation was blocked by heat inactivation or by pepstatin but not with other protease inhibitors (Fig. S11), suggesting that degradation was due to aspartic proteases. Accordingly, two (Pep1, Pep2) of the four major reported acidic proteases (Pep1, Pep2, CtsD and OpsB) were identified in the conidial cell wall (Reichard et al., 1997; Sriranganadane et al., 2010) (Table 1) and in the filtrate of germinating conidia (O. Kniemeyer, personal communication). In agreement with these results, recombinant Pep1 and Pep2 proteases were indeed able to degrade native RodA (Fig. 10).

Therefore, to understand the role of these aspartic proteases on the degradation of rodlet layer during germination, a quadruple mutant of *A. fumigatus*, devoid of the four major aspartic proteases ($\Delta pep1/\Delta pep2/\Delta ctsD/\Delta opsB$), was constructed (Figs. S14-S17). Surprisingly, neither the degradation of RodA nor the germination of the quadruple mutant were affected (data not shown), indicating that even though these major aspartic proteases degrade RodA, other proteases can degrade the rodlet layer during conidial germination.

Discussion

The structure and assembly of A. fumigatus RodA into rodlets

Analysis of rRodA and conidial native RodA for understanding of the assembly of the rodlets has suggested that rodlet formation in *A. fumigatus* may be unusual in that the two amyloidogenic regions are likely to be involved in the formation of intermolecular β -sheet structures. In contrast in other fungi, only single amyloidogenic regions have been identified (Macindoe et al., 2012; Niu et al., 2017; Wang et al., 2004). Here, like earlier studies with the *Neurospora crassa* hydrophobin EAS (Kwan et al., 2006), we used solution NMR spectroscopy to determine the structure of the monomeric form. The RodA monomer in solution displays the β -barrel structure typical of hydrophobins and the surface of RodA exhibits a separation of charged and hydrophobic amino acid residues, which renders the protein amphipathic. Also, as for other hydrophobins, the amphiphilic nature of the RodA monomer likely plays an important role in localizing the protein at an interface and the flexibility of the amyloidogenic regions may be necessary to accommodate the structural changes required for multimerization and uniform covering of substrates. In the absence of crystals of hydrophobins in the assembled form, solid state NMR (ssNMR) has been used to probe the 3D structure of the EAS rodlets and confirmed that hydrophobin rodlets are composed of a β -sheet-rich structure with an altered protein fold compared to the monomer in solution (Morris et al., 2012). Similar ssNMR studies of RodA will likely reveal how the two amyloidogenic regions in the RodA rodlets contribute to the structure and packing of rodlets within the robust protein film.

Earlier studies have shown that an increase in pH, the presence of Ca^{2+} , a low ionic strength, the polarity of the solvent, the high temperature, the presence of detergents, the protein concentration, specific nutrients and the physico-chemical nature of the surface are all environmental factors which can influence rodlet formation (Aimanianda et al., 2009; Cicatiello et al., 2017; Longobardi et al., 2012; Morris et al., 2011; Niu et al., 2017; Pham et al., 2016; Talbot et al., 1993, 1996; Tanaka et al., 2017; Wang et al., 2004; Zykwiniska et al., 2014a). The comparison of rodlet formation at the surface of the conidia of *A. fumigatus* and on an artificial hydrophobic surface like HOPG of wild-type or rRodA mutated proteins, has indicated that the environment influences the formation of the rodlet structure. For example,

even though rRodA carrying mutations form rodlets that are similar to the wild type protein on the HOPG substrate, the rodlet structure and coverage on the conidial surface of the mutant looked altered on the more complex surface of the conidial cell wall. Similar observations have been reported for other hydrophobins like SC3 from *Schizophyllum commune* and Vmh2 from *Pleurotus ostreatus* that can also spontaneously self-assemble in aqueous solutions (Longobardi et al., 2012; Zykwiniska et al., 2014a). The presence of oligosaccharides facilitates the self-assembly in water of *P. ostreatus* Vmh2 and the width of the rodlet is influenced by the addition of cyclic 1,4-glucans (Armenante et al., 2010). The transition of *Schizophyllum commune* SC3 from a helical configuration to the stable β -rich amyloid conformation is promoted by the presence of cell wall polysaccharides (Scholtmeijer et al., 2009). In *A. fumigatus*, the role of the inner components of the conidial cell wall, which are located below the rodlet layer, such as α - and β -1,3-glucans, galactomannan or melanin, on the structure and anchoring of rodlets has not been investigated to date. An accurate definition of the events controlling the structural conversion of the *A. fumigatus* RodA protein into rodlets remains to be generated and should take into account the natural environment of the conidium. Moreover, the observation that there are different triggers for rodlet assembly in different hydrophobins likely reflects the fact that apart from the pattern of conserved disulfide bonds, which constrains the β -barrel core, the sequences and structures of the proteins are highly diverse (Ren et al., 2013).

In vitro studies, while very useful, can be limited by the homogeneous nature of the substrate and the inability to effectively probe the three-dimensional organization of the rodlet-containing layer. Moreover, the role of the cellular environment used in the heterologous host for the production of recombinant RodA protein has not been investigated. Even though *A. fumigatus* RodA protein has been produced for our study in *E. coli* and folded *in vitro* to obtain a protein that is folded like conidial RodA and is able to form rodlets, it has been previously shown that *A. fumigatus* RodA protein can be also produced in *Pichia pastoris* (Pedersen et al., 2011). The amount produced in *P. pastoris* was 200 to 300 mg/L to be compared with *E. coli* (~2 mg/L). However, although it was verified that the recombinant RodA of *A. fumigatus* produced in *Pichia* was able to convert a hydrophilic substrate into a hydrophobic surface, the ability of the recombinant proteins produced in *P. pastoris* to make rodlets and their glycosylation were not tested.

On conidia the rodlets are packed into distinct nanodomains, with a width of 60 to 100 nm while the height of the layer is estimated to be around 5 nm (this study and (Zykwiniska et al., 2014b)), suggesting that the nanodomains are organized in multilayers on the surface of the conidium. The parallel organization of rodlets could arise from the assembly of the oligomers directly on the conidium surface or from the adhesion of rodlets aggregated in the medium prior to binding to the cell wall. The growth of one rodlet may provide a surface that favours nucleation and elongation of an adjacent rodlet, a process of secondary nucleation that has been shown to be important in the formation of some pathogenic amyloids (Pham et al., 2016). The observation that rodlets in adjacent nanodomains are oriented at angles of up to 90° may be consistent with repulsion between the outer hydrophobic components of different layers but could also be a consequence of the termination of nucleation and/or elongation when coverage of the conidium surface by rodlets removes the available surface. In addition, if acid extraction leads to the production of a RodA band lacking 40 aminoacids (Fig. S7) the incubation with hydrofluoric or formic acid did not degrade recombinant RodA to form the protein lacking N-terminal 40-amino acids. These results suggest that the aminoacid 40 or 41 is a weak point in the rodlets structuration, which may be involved in the binding of RodA to the inner cell wall layer.

While RodA clearly contributes to conidial hydrophobicity, an additional element to take into account is the role of other members of the hydrophobin family in controlling the hydrophobic properties of the conidium. In *A. nidulans*, all six hydrophobins contributed to hydrophobicity of the spore surface and were able to self-assemble (Grünbacher et al., 2014). However, only the deletion of *RODA* led to a mutant without rodlet on the surface even though DewA was able to form rodlets *in vitro*. When DewA and DewB were placed under the control of the *RODA* promoter in a $\Delta rodA$ mutant, both hydrophobins were able to produce rodlets, albeit with a different structure than the classical parental strain. Similarly, when the hydrophobins of *S. commune* (SC1 and 4) or RodA and Dew A of *A. nidulans*, or EAS of *N. crassa* were expressed from the *mpg1* promoter of *M. oryzae*, they were also able to complement, although only partially, the *mpg1* mutation, showing at the same time the lack of specificity of the hydrophobin aggregation but the need to recognize specific fungal features to induce the proper formation of rodlets on the conidial surface (Scholtmeijer et al., 2009; Kershaw et al., 1998). In *Beauveria bassiana*, the single hydrophobin Hyd1 is able to form rodlets although the rodlets are truncated and incomplete suggesting that the two hydrophobins Hyd1 and 2 of this species act together to form the rodlets. In contrast in *A. fumigatus*, although eight hydrophobin genes (*RODA* to *G*) were identified (Jensen et al., 2010), only RodA was responsible for the formation of rodlets since a septuple *RODB/C/D/E/F/G* mutant with all these genes deleted but with wild-type RodA appeared like the parental strain (Valsecchi et al., 2017). The function of the other hydrophobins in *A. fumigatus* remains unknown.

Role of rodlets in fungal pathogenicity

Several biological reasons could explain the role of rodlets in the pathogenicity of *A. fumigatus*. First, the RodA protein itself, even in drastically different conformations or with point mutations is immunologically inert. The reasons for this lack of immune activation remain unknown and unlike other functional and disease-associated β -amyloids, which stimulate the innate cells via TLR and inflammasome activation (Halle et al., 2008; Tükel et al., 2009), the structure of hydrophobin proteins does not dictate immune inertness. Our data on RodA and recent unpublished data (Heddergott and Latgé) on other cysteine-rich proteins of low Mr suggests that the amino acid sequence is responsible for the lack of immunogenicity against RodA. Second, by masking inner cell wall PAMPs like β -glucans and mannans, the rodlets are responsible for the lack of recognition of conidia by the immune cell pattern recognition receptors (Bayry et al., 2014; Beauvais et al., 2013). In a murine model of corneal infection, the $\Delta rodA$ mutant is less pathogenic because of an increase in cytokine production, which boosts the anti-fungal immune response (Carrion et al., 2013). Third, conidial rodlets have a negative impact on the innate immune mechanisms: they reduce NET formation when neutrophils come into contact with conidia and the addition of RodA to $\Delta rodA$ mutant conidia reduces the formation of NET (Bruns et al., 2010) and decrease macrophage efficacy (Dagenais et al., 2010). Fourth, the rodlet layer makes the conidium less permeable to the external antifungal molecules. The loss of rodlets and change in hydrophobicity that occurs during germination is indeed associated with an increased sensitivity to antifungal drugs (Dague et al., 2008a). Finally, hydrophobins favour the binding of enzymes to their substrate and stimulate their enzymatic activity (Pham et al., 2016; Ribitsch et al., 2015; Takahashi et al., 2005, 2005; Tanaka et al., 2017). This has been especially shown with MPG1 and two orthologs of RodA, namely *A. oryzae* RolA and *A. nidulans* RodA that have been shown to interact via ionic interaction with enzymes such as cutinases. It remains to be investigated whether fungal enzymes involved in pulmonary matrix

degradation and host collectins and antimicrobial peptides bind to hydrophobin rodlets and play a role during lung infection by *A. fumigatus*.

Conclusions

The hydrophobin, RodA, forms functional amyloids with a rodlet morphology that coat *A. fumigatus* conidia. Assembly of parental and point mutated RodA proteins *in vitro* and on the conidial surface was analyzed with multiple technical approaches. Proper rodlet assembly requires the presence of intact cysteine bridges and two amyloidogenic regions. Only an intact rodlet layer is able to hide the conidia whereas even a partial rodlet disorganization can activate the host immune system. Degradation of the rodlet layer during germination by the fungal cell wall associated aspartic proteases could also activate immune cells. These studies have shown that the unique structure of the rodlets on the surface of the aerial conidia of human fungal pathogens plays a major role during the interaction with the host. It is a dynamic structure that continuously evolves from conidial formation to germination. Progress in understanding the mechanisms of self-assembly and the rodlet structure may lead to advances towards solving medical problems associated with amyloid aggregation.

Material and methods

Extraction of the native RodA from conidia of *A. fumigatus*

RodA was extracted from dry conidia of *A. fumigatus* by a 2h incubation on ice with formic acid. The material extracted was similar to the one extracted with a 48% hydrofluoric acid extraction for 72 h at 4°C shown before to isolate rodlets from conidia (Aimanianda et al., 2009). The presence of rodlets was analyzed by SDS-PAGE using a 15% polyacrylamide gel and by Western blotting with an anti-RodA antibody (Valsecchi et al., 2017).

Production of the recombinant RodA protein for NMR analysis

The sequence used to prepare the recombinant RodA protein for solution NMR corresponds to residues 19-159 of *A. fumigatus* A1163 RodA (AFUB_057130) without the predicted N-terminal secretion peptide and containing an extra serine N-terminal residue that arises from cloning. The protein was obtained in lyophilized form as described (Pille et al., 2015) with a slight modification. Briefly, the HisTag-ubiquitin-rRodA fusion protein was encoded on a pET-28b(+) plasmid (Proteogenix) conserving the C-terminal ubiquitin GG recognition motif for deubiquitinases. The proteolysis step to separate RodA from the hexa-histidine-tagged ubiquitin was performed with His-tagged-deubiquitinase UBP41. Uniformly ^{15}N and $^{15}\text{N}/^{13}\text{C}$ doubly labeled rRodA was expressed in minimal media containing $^{15}\text{NH}_4\text{Cl}$ and $\text{U-}^{13}\text{C}_6$ glucose (Euriso-top, Saclay) as sole sources of nitrogen and carbon, respectively. The rRodA was obtained after *in vitro* oxidative refolding. Integrity, label and sequence of the protein were analyzed by mass spectrometry. Protein samples were typically prepared at 0.36-0.69 mM concentrations in 20 mM CD_3COONa pH 4.3 10% D_2O . No dependence of line width or chemical shifts, and no signal intensity variation over time were observed in $^1\text{H-}^{15}\text{N}$ HSQC between 0.10 and 0.69 mM concentrations, indicating that the soluble form does not self-assemble in this concentration range and that samples were stable.

Doubly labeled conidial RodA was extracted from *A. fumigatus* conidia grown in minimal media supplemented with $^{15}\text{NH}_4\text{Cl}$ and $\text{U-}^{13}\text{C}_6$ glucose. Mature conidia were harvested and washed extensively in Tween 20 0.05%. To recover RodA protein, lyophilized conidia were incubated with 100% formic acid on ice (10 min to 2 h). After removal of the conidia by centrifugation, the supernatant was flushed with gaseous nitrogen to remove formic acid and the protein was then dissolved in water. NMR samples were prepared at 50-60 μM concentrations in the same buffer that was used for recombinant RodA.

NMR and structure calculation methods

Experiments were performed on a 14.7 T Direct Drive 600 NMR System spectrometer (Agilent Technologies, Santa Clara) equipped with a triple resonance cryogenic probe. Spectra were recorded at 25°C using VnmrJ 3.2A (Agilent). Processing and analysis were performed with NMRPipe and CCPNMR Analysis, respectively. Proton, carbon and nitrogen resonances were assigned to backbone and side-chains atoms following a standard strategy with 2- (2D) and 3-dimensional (3D) heteronuclear experiments as described by Pille and co-workers (Pille et al., 2015). Assignment of RodA extracted from conidia was performed by comparing ^1H - ^{15}N and ^1H - ^{13}C HSQC, HNCQ, HNCA experiments with the corresponding spectra of rRodA. Backbone amide ^{15}N heteronuclear ^1H - ^{15}N nOe (nuclear Overhauser enhancement) spectra were recorded to analyze the internal dynamics of rRodA, with a saturation/no-saturation delay of 3 s and a total recovery delay of 3.07 s. To obtain distance constraints, 3D ^{15}N and ^{13}C (aliphatic and aromatic regions) edited HSQC-nOesy were recorded with a 120 ms mixing time for nOe build-up (Kay et al., 1993; Muhandiram and Kay, 1994). Dihedral Phi and Psi angle constraints were calculated with Talos-N (Shen et al., 2009). An HNHA spectrum was used to validate Talos-N Phi angle constraints. Assignment of nOes and structure calculations were performed using ARIA 2.3.2 (Rieping et al., 2007) coupled to CNS 1.2.1 (Brunger et al., 1998). ARIA was run with a log-harmonic potential and automatic weighting of constraints spin diffusion correction, network anchoring and explicit water refinement using standard protocols. The ten lowest total-energy conformers were selected out of the 500 structures calculated in the final run. Structures were calculated with the disulfide bridge pairing determined experimentally for native EAS (Kwan et al., 2006) and observed in other recombinant hydrophobins [C1-C5 (56-133), C2-C6 (64-127), C3-C4 (65-105), C7-C8 (134-152)]. Herein, the cysteines are noted by their order of appearance in the sequence (C1 to C8) or by their sequence position. A careful analysis of the topology of the molecule (antiparallel β -sheets involving cysteines), of the network of nOes supporting the topology and the unambiguous assignment of several nOes between cysteines and involving neighbouring residues clearly demonstrated that rRodA showed the topology determined for native EAS. Structures were visualized and analyzed with PYMOL (Schrödinger LLC). Secondary structure content of RodA was determined according to DSSP. The quality of the structures was assessed with PROCHECK 3.5.4 (Laskowski et al., 1993), WHATCHECK (Hoofst et al., 1996), MOLPROBITY (Chen et al., 2010) and PROSA (Sippl, 1993). The coordinates have been deposited to the pdb under the accession code 6GCJ.

Production of the recombinant RodA and mutant proteins for Thioflavin T (ThT) fluorescence and AFM imaging

Production of rRodA proteins, wild type and mutants, was achieved as previously described for the hydrophobin MPG1 (Pham et al., 2016) apart from the following three changes. Fusion proteins were expressed using a modified version of the pHUE plasmid, encoding a version of ubiquitin with the C-terminal GG residues of ubiquitin replaced by a TEV protease recognition sequence. Oxidative refolding was achieved by dialysis against 10 mM reduced glutathione, 1 mM oxidized glutathione, 50 mM sodium acetate, 100 mM sodium chloride, pH 5.0) with two changes of redox buffer (6 h each at 4°C; protein:buffer 1:25 (v/v)). Cleavage of hydrophobin from His₆-ubiquitin occurred by addition of TEV enzyme. After reverse phase HPLC, the correct folding of rRodA and variants was confirmed by 1D ^1H nuclear magnetic resonance spectroscopy.

Thioflavin T (ThT) rodlet self-assembly assay

Lyophilized rRodA and mutant proteins were first solubilized in deionized H₂O. The stock protein solutions were diluted in ThT buffer (40 μM Thioflavin T in 20 mM sodium acetate,

pH 5.0) to a final protein concentration of 25 $\mu\text{g}/\text{mL}$ (1.73 μM). Samples were prepared to 100 μL in a black, clear-bottom 96-well microplate (Greiner Bio-One). The microplate was incubated at 50°C and fluorescence recorded with 440/480 nm excitation/emission filters following every 90 s of double orbital shaking at 700 rpm in a BMG Labtech POLARstar Omega multi-mode microplate reader BMG Labtech Australia, VIC, Australia. All protein samples in each independent experiment were analyzed in triplicate. Each sample triplicate was normalized and graphed individually. Lag phase and time to reach half of the maximum fluorescence ($t_{1/2}$) was interpreted from mean of triplicates. Lag phase and $t_{1/2}$ were determined by the last time point before the normalized ThT fluorescence reached a value of 0.1 and the time at which normalized 0.5 ThT fluorescence reached 0.5, respectively. Statistical analysis was performed on GraphPad Prism (version 6.04 for Windows, GraphPad Software, CA, USA, www.graphpad.com). Data was evaluated using one-way ANOVA followed by Tukey's multiple comparisons test.

Surface structure of in vitro produced recombinant proteins and peptides seen with atomic force microscopy or transmission electron microscopy

Protein samples were diluted to 5 $\mu\text{g}/\text{mL}$ (0.35 μM) in deionized H_2O . For rRodA-WT and point mutations, a 50 μL drop of protein was deposited onto a freshly-cleaved highly-ordered pyrolytic graphite (HOPG) surface (Holgate Scientific, NSW, Australia) and allowed to dry down in a covered, ambient chamber overnight at room temperature. rRodA mutant proteins containing multiple mutations were prepared similarly, but were incubated at 50°C in a humid environment for 2 h, followed by overnight drying in the ambient chamber. The HOPG surfaces were then further dried in a 70°C oven for 2 h to remove any excess moisture.

The morphology of the assembled protein layers on the HOPG surface was imaged at ambient temperature using a Multimode Nanoscope[®] III atomic force microscopy (Veeco, CA, USA). The HOPG surfaces were scanned with tapping mode using a silicon scanning probe with tip radius < 10 nm, force constant of 40 N/m and resonance frequency of 300 kHz (Tap300AI-G, BudgetSensors[™], Bulgaria). The images were further processed with Gwyddion software (version 2.38, <http://gwyddion.net>).

For TEM, a 20- μL droplet of peptide (200 $\mu\text{g}/\text{mL}$) was placed on a formvar/carbon coated copper TEM grid (200-mesh, ProSciTech), and incubated for 2 min at room temperature. Excess solution was removed with filter paper and grids were washed twice by floating the grids on water droplets and wicking away the excess solution. Samples were stained with 2% uranyl acetate and imaged on a JEOL 1400 JEOL TEM equipped with a Morada G3 16 megapixel side camera. All images were analyzed with ImageJ.

Construction of the RodA point mutated and N-terminus minus ΔrodA mutants

Construction of the template plasmid for mutagenesis

The template for directed mutagenesis reactions was constructed with the pUC19 plasmid containing the RodA upstream region (1486 bp), the RodA open reading frame (583 bp), the hygromycin cassette beta recombinase (4767 bp) (Hartmann et al., 2010) and the RodA downstream region (853 bp). The pUC19 plasmid was digested with *SacI* and *PstI* and the hygromycin cassette was excised with *FspI* from plasmid pSK529 (Hartmann et al., 2010) and the corresponding band was purified from an agarose-gel. The RodA upstream region, open reading frame and downstream region were amplified by PCR using as template the gDNA of the CEA17_ $\Delta\text{akuB}^{\text{KU80}}$ strain (da Silva Ferreira et al., 2006) using primers 1, 2, 3, 4 (see Supplementary Table 3). The final template plasmid was assembled with the GeneArt Seamless Cloning and Assembly kit (Thermo Fisher).

Construction of the point mutated RodA and N terminus minus strains

RodA point mutants were obtained using the QuikChange Multi Site-Directed Mutagenesis Kit (Agilent technologies) following the manufacturer instructions. Each PCR-mutagenesis reaction was performed with the template plasmid described above and primers as described in Supplementary Table 3. PCR-mutagenesis reactions for mutations C127S, C64S/C65S/C133S/C134S, I114G, I115S, L145S, I146G and D140G were performed with the template plasmid described above and primers 5, 6-7, 8, 9, 10, 11 and 12 (Supplementary Table 3), respectively. The plasmid for the double mutation I115S/I146G was generated with the RodA I146G plasmid and primer 9. DNA sequencing was performed after each mutagenesis reaction to verify the presence of the expected mutation. Each mutated plasmid was digested with the *SphI* restriction enzyme before electroporation on the CEA17_Δ*akuB*^{KU80} parental strain as described by Lambou *et al.* (2006). Transformants were selected with 150 μg/ml of hygromycin. PCR and Southern blot analysis were performed to verify the transformants. In the Southern blot analysis, the gDNAs of mutated strains were digested with *BstZ17I* and annealed with a PCR-DIG labeled probe (Roche) of the RodA ORF (Fig. S5). A PCR was performed using primers that anneal outside the integration site of the cassette for DNA sequencing in order to verify the frame, the point mutations and the presence of the 120 bp remaining after the removal of the *HPH*^R-β*rec* cassette.

The RodA-N terminal cassette was constructed with four DNA fragments, I, II, III, IV described in Fig. S4. Fragments I, II and IV were amplified by PCR with the following pairs of primers 1-13, 14-2 and 3-4 (see Supplementary Table 3). Fragment I and II contain the *RODA* upstream region and the *RODA* ORF except the DNA bases from C55 to C112 corresponding to the aminoacids P20-V39. Fragment III, corresponding to the *HRP*^R-β*rec* resistance, was digested with *FspI* from plasmid pSK485 followed by gel band purification. Fragment IV contains the *RODA* downstream region. The 4 fragments were assembled each other by GeneArt Seamless Cloning and Assembly kit (Thermo Fisher) and were used for electroporation after digestion with *SphI*, on the CEA17 Δ*akuB*^{KU80} parental strain (da Silva Ferreira *et al.*, 2006).

Complementation of RodA I146G and C64S/C65S/C133S/C134S strains

The same plasmid constructed as a template for the mutagenesis reaction was used to complement the RodA I146G and C64S/C65S/C133S/C134S strains. This plasmid was digested with *SphI* before transformation by electroporation (Lambou *et al.*, 2010).

Construction of quadruple Δ*ctsD*/Δ*opsB*/Δ*pep1*/Δ*pep2* mutant strain

In order to produce the quadruple aspartic protease mutant, four different deletion cassettes were constructed to delete the four major genes coding for aspartic acid proteases (*CtsD*, *OpsB*, *Pep1* and *Pep2*) of *A. fumigatus* (Monod, 2008) and to obtain the Δ*ctsD*, Δ*opsB*, Δ*pep1* and Δ*pep2* mutants. Each cassette consists in three DNA fragments, the upstream and the downstream DNA regions of the specific gene and the Hygromycin β *rec* resistance (*HPR*^R-β*rec*) (Part A of Fig. S12 to S15). Upstream and downstream regions were amplified by PCR using the pair of primers listed in Supplementary Table 3: Δ*ctsD* (15-16, 17-18), Δ*opsB* (19-20, 21-22), Δ*pep1* (23-24, 25-26) and Δ*pep2* (27-28, 29-30). The *HPR*^R-β *rec* was digested with *FspI* from plasmid pSK485 followed by gel band purification. The three DNA fragments of each cassette were assembled with the GeneArt Seamless Cloning and Assembly kit (ThermoFisher). Transformation by electroporation was done as described (Lambou *et al.*, 2010). Integration of the cassette and deletion of the targeted gene at the right locus was verified by Southern blot as described in the Part B of Fig. S12 to S15). To construct the

quadruple mutant, the $\Delta cstD::HPH^R-\beta rec$ mutant was first constructed. After the verification of the deletion at the right locus and the presence of only one single insert (see above) $\Delta cstD$ mutants were grown on MM-glc+2% xylose to excise the hygromycin cassette by the β recombinase through the six sites-specific (Hartmann et al., 2010). Then the $\Delta ctsD::six/\Delta opsB::HPH^R-\beta rec$ double mutant was constructed as described above with the plasmid shown in Fig. S14. Excision of the $HPH^R-\beta rec$ marker as well as the third and fourth deletions was as described above to produce a $\Delta ctsD::six \Delta opsB::six \Delta pep1::six \Delta pep2$ quadruple mutant.

Growth conditions of the A. fumigatus strains and analysis of the conidial phenotype of the mutants.

All the strains were maintained on 2% malt agar slants at room temperature. Conidia were recovered from these slants using 0.05% Tween20 aqueous mixtures and conidial germination was followed microscopically and quantified on 2% Malt agar medium. Survival of the conidia in presence of SDS (0.01 to 0.1%) and H_2O_2 (0.5 to 3 mM) was also tested to evaluate cell wall permeability changes.

Atomic force microscopy of conidia

Conidial surfaces were analyzed by AFM, using a Multimode VIII AFM (Bruker, Santa Barbara, CA). *A. fumigatus* conidia were immobilized by mechanically trapping them into porous polycarbonate membranes with a pore size similar to the conidium diameter (it4ip SA, Belgium). After filtering a concentrated suspension of conidia, the filter was rinsed with deionized water, carefully cut and attached to a metallic puck using double-sided sticky tape. The mounted sample was then transferred to the AFM liquid cell. Images were performed in contact mode under minimal applied force, using oxide-sharpened microfabricated Si_3N_4 tips (MSCT, Bruker) with a nominal spring constant of 0.01 N/m. The surface coverage of the amorphous layer and the rodlets were determined visually using ImageJ software.

Analysis of the immune response against recombinant RodA proteins and conidia of the RodA mutants

Recombinant proteins used for studying the immune response

During the reverse phase chromatography purification step of rRodA, two main peaks corresponding to RodA are observed. The peak with lower retention time upon elution with a gradient of increasing acetonitrile in 10% methanol 0.01 %TFA) corresponded to the folded protein as determined by proton 1D and $^1H-^{15}N$ HSQC NMR spectra. The protein eluting with a higher retention time, either alone or with the reducing agent TCEP (Tris(2-carboxyethyl)phosphine, 5 mM) showed the characteristic NMR spectra of an unfolded protein. To produce reduced and cysteine blocked RodA samples, lyophilized, unfolded rRodA was dissolved in 6 M guanidinium hydrochloride-Tris-HCl 100 mM pH 7 or 8. Potential disulfides were reduced by incubating rRodA with DTT (dithiothreitol) at 1:10 cysteine/DTT ratio at 37 °C under a nitrogen flow. Reduced cysteines were then alkylated by adding a ten-fold molar excess of alkylating agent relative to DTT at pH 7 with iodoacetamide or at pH 8 with N-ethyl maleimide and incubating the sample 1 h at 37 °C under a nitrogen flow. Alkylation was stopped with a ten-fold molar excess of β -mercaptoethanol relative to the alkylating agent. Alkylated rRodA samples were purified by reverse phase chromatography like rRodA, lyophilized and kept at 4 °C. SELDI (surface enhanced laser desorption ionization time of flight) mass spectrometry indicated a complete labelling for IA, and two species with 7 or 8 alkylated cysteines for NEM.

Recombinant RodA wild type and proteins carrying mutations were produced as described

above. Conidia from parental and mutant strains were produced after 3, 10 or 30 days of culture on 2% malt agar. Conidia were recovered from slants by vortexing the culture with an aqueous 0.05% Tween 20 solution. Conidia were extensively washed with water. Surface material of the mutant conidia was extracted after incubation of the conidia for 2 h in formic acid. The material was dialyzed before being tested immunologically.

Human material for immune analysis

PE-conjugated MAbs to CD83 (clone HB15e), CD127 (clone HIL-7R-M21); FITC-conjugated MAbs to CD86 (clone 2331 (FUN-1)), CD25 (clone M-A251); Alexa 700-conjugated MAbs to CD4 (clone RPA-T4) were from BD Biosciences. APC-conjugated MAbs to FoxP3 (clone 236A/E7) and Fixable Viability Dye eFluor 506 were from eBioscience.

Peripheral blood mononuclear cells (PBMCs) were obtained from buffy bags of healthy donors by Ficoll density gradient centrifugation. Anonymized buffy bags of the healthy blood donors were purchased from Centre Necker-Cabanel, Etablissement Français du Sang, Paris, France. Ethical committee permission was obtained for the use of buffy bags of healthy donors (Institut National de la Santé et de la Recherche-EFS ethical committee convention 15/EFS/012). Monocytes were isolated from PBMCs by positive selection using the human CD14. Monocytes (0.5×10^6 cells per ml) were cultured in the presence of granulocyte-macrophage colony-stimulating factor (GM-CSF; 1,000 IU per 10^6 cells) and IL-4 (500 IU per 10^6 cells) (both cytokines from Miltenyi Biotec) for 5 days to obtain immature monocyte-derived DCs.

Autologous naïve T cells were negatively isolated in a two-step process. First untouched CD4 T cells were isolated using the CD4 T cell isolation kit II. In the second step CD4 T cells were labeled with CD45RO and CD25 and the negative fraction containing the $CD4^+CD45RA^+CD25^-$ was used for assessing naïve T cell responses. All the isolation kits were purchased from Miltenyi Biotec (Paris, France).

Treatment of DCs with RodA proteins and conidia

Immature DCs (0.5×10^6 /ml) were cultured in the presence of GM-CSF and IL-4 alone or with parental or mutant conidia (1:1) for 48 h. In other experiments, DCs were incubated with various forms of RodA ($1 \mu\text{g}/0.5 \times 10^6$ DCs/ml) or LPS ($100 \text{ ng}/0.5 \times 10^6$ DCs/ml). After 48 h of incubation, cell-free culture supernatants were collected for analysis of IL-6 and TNF α DC-cytokines (ELISA Ready-SET-Go, eBioscience). DCs were processed for flow cytometry by surface staining with fluorochrome-conjugated antibodies. Data were acquired on a LSR II (BD Biosciences) flow cytometer and analyzed using the BD FACS DIVA software (BD Biosciences).

DCs-naïve CD4⁺ T cell co-culture

To explore the effect of *A. fumigatus* conidia-stimulated DCs on CD4⁺ T cell polarization, after stimulation, DCs were extensively washed and co-cultured with 0.1×10^6 $CD4^+CD45RA^+CD25^-$ autologous naïve T cells at a 1:20 ratio for five days in serum-free X-VIVO medium. Cell-free supernatants were collected after 5 days. Surface staining of T cells was performed with fluorescence-conjugated mAbs against CD4, CD25 or CD127. Then, cells were fixed, permeabilized using an intracellular staining kit (eBioscience) and incubated at 4 °C with fluorochrome-conjugated mAbs against FoxP3. Data were recorded on a LSR II (BD Biosciences) flow cytometer and analyzed with the BD FACS DIVA software (BD

Biosciences). The cell-free culture supernatants were collected and analyzed for various T cell cytokines by ELISA (IFN- γ , IL-4, and IL-17A; ELISA Ready-SET-Go, eBioscience).

Proteolytic degradation of RodA

In a total volume of 30 μ l containing 20 mM acetate buffer pH 5.5, native RodA (5 μ g) recovered from conidia with hydrofluoric acid was incubated for 2 h with an ethanol-precipitated protein mixture from the culture supernatant (2 μ g of protein) of *A. fumigatus* grown in Brian's medium at 37 °C for 24 h. The nature of the proteases degrading RodA was investigated by addition of different protease inhibitors at 1 mM concentration. The role of aspartic acid proteases was confirmed by incubating conidial RodA with the recombinant aspartic acid proteases rPep1 or rPep2 at 37 °C for 2 h (Sarfati et al., 2006): RodA (4 μ g) + rPep1 or rPep2 (1 μ g) in 30 μ l of 20 mM acetate buffer, pH 5.5. Degradation of RodA was monitored by SDS-PAGE.

Mass spectrometry experiments

In-solution digestion

Conidia were incubated in formic acid for 2 hours as described above. Preliminary experiments showed that the same proteins were extracted by 10 min or 2 hours incubation times. Dried samples were re-suspended in 100 μ L 8 M urea, 100 mM Tris-HCl pH 8.5. Briefly, samples were reduced with 5 mM TCEP for 30 minutes and alkylated with 10 mM iodoacetamide for 30 minutes at room temperature in the dark. Protein samples were then incubated with 250 ng rLys-C Mass Spec Grade (Promega, Madison, WI, USA) for 5 hours at 37°C for the first digestion. Samples were then diluted to 2 M urea with 100 mM Tris HCl pH 8.5 and 500 ng sequencing grade modified trypsin (Promega, Madison, WI, USA) was added for the second digestion overnight at 37°C. A second incubation with the same amount of trypsin (5 hours at 37°C) was performed to ensure a complete digestion. Digestion was stopped by adding formic acid and peptides were desalted and concentrated on a Sep-Pak C₁₈ SPE cartridge (Waters, Milford, MA, USA) according to manufacturer's instructions.

Mass spectrometry analysis

Tryptic peptides were analyzed on a Q Exactive Plus instrument (Thermo Fisher Scientific, Bremen) coupled with an EASY nLC 1000 chromatography system (Thermo Fisher Scientific). Sample was loaded on an in-house packed 50 cm nano-HPLC column (75 μ m inner diameter) with C₁₈ resin (1.9 μ m particles, 100 Å pore size, Reprosil-Pur Basic C18-HD resin, Dr. Maisch GmbH, Ammerbuch-Entringen, Germany) and equilibrated in 98 % solvent A (H₂O, 0.1 % FA) and 2 % solvent B (ACN, 0.1 % FA). Peptides were first eluted using a 2 to 18 % gradient of solvent B during 112 min, then a 18 to 30 % gradient of solvent B during 35 min, a 30 to 45 % gradient of solvent B during 15 min and finally a 45 to 60 % gradient of solvent B during 5 min all at a 250 nL/min flow rate. The instrument method for the Q Exactive Plus was set up in the data dependent acquisition (DDA) mode. After a survey scan in the Orbitrap (resolution 70 000), the 10 most intense precursor ions were selected for HCD fragmentation with a normalized collision energy set up to 27. Charge state screening was enabled, and precursors with unknown charge state or a charge state of 1 and > 7 were excluded. Dynamic exclusion was enabled for 45 s.

Data processing for protein identification and quantification

All data were searched using Andromeda (Cox et al., 2011) with MaxQuant software (Cox and Mann, 2008; Tyanova et al., 2016) version 1.5.3.8 against a *Neosartorya fumigata* (strain CEA10 CBS 144.89 FGSC A1163) (*Aspergillus fumigatus*) Uniprot database (Proteome ID

UP000001699, 9942 entries) concatenated with mutated RodA protein, usual known mass spectrometry contaminants and reversed sequences of all entries. Andromeda searches were performed choosing trypsin as specific enzyme with a maximum number of two missed cleavages. Possible modifications included carbamidomethylation (Cys, fixed), oxidation (Met, variable) and Nter acetylation (variable). The mass tolerance in MS was set to 20 ppm for the first search then 6 ppm for the main search and 10 ppm for the MS/MS. Maximum peptide charge was set to seven and five amino acids were required as minimum peptide length. The “match between runs” feature was applied for samples having the same experimental condition with a maximal retention time window of 0.7 minutes. One unique peptide to the protein group was required for the protein identification. A false discovery rate (FDR) cut-off of 1 % was applied at the peptide and protein levels. Reverse proteins and usual MS contaminants were removed before the analysis of the data. Quantification of each identified protein was performed by summing the intensities of its associated peptides.

Analysis of protein intensity data

For the quantification of the proteins and comparison of conditions, proteins with at least two quantified intensity values among the replicates of the condition were selected for further analysis.

Selection of “condition-specific proteins” was made by the selection of proteins with peptides identified in a unique condition. For ranking these condition-specific proteins from the most abundant ones to the least in the condition where they are observed, we used the average of their observed iBAQ values among the replicates of the condition. The iBAQ algorithm consists to normalize the summed peptide intensities by the number of theoretically observable peptides for each protein and is a relevant method to rank the absolute abundance of different proteins within a single sample (Schwanhäusser et al., 2011). Since we were particularly interested in secretory proteins among all these condition-specific proteins, we used the SignalP algorithm to predict proteins with secretory signal peptides (Petersen et al., 2011). This algorithm was used with the default cut-off value, which optimizes the performance measured as Matthews Correlation Coefficient (MCC). The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Table 1) partner repository with the dataset identifier PXD008503.

Funding

This work was supported partially by the grant Aspergillus by Aviesan, the grant DEQ20150331722 LATGE Equipe FRM 2015 and the French Australian FAST project FR110012. This work was additionally funded by the French ANR (HYDROPHOBIN ANR-CR-16115-01 and FUNHYDRO ANR-16S-CE110020-01) and the Institut Pasteur PTR 529. AP was a recipient of an Université Pierre et Marie Curie (UPMC) fellowship. The NMR spectrometer was partially supported by the Région Ile de France (SESAME grant).

The authors declare no conflict of interest
JP Latgé

Acknowledgements

The authors thank Olaf Kniemeyer (HKI, Jena, Germany) for providing unpublished MS data of the secretome of *A. fumigatus*, Remi Beau and Anne Beauvais (Aspergillus unit, Institut Pasteur) for providing the anti-RodA polyclonal antibody, Prof. Chiaro Neto for use of the AFM and the staff of Sydney Microscopy and Microanalysis for maintenance of the TEM.

References

- Aimanianda, V., Bayry, J., Bozza, S., Knemeyer, O., Perruccio, K., Elluru, S.R., Clavaud, C., Paris, S., Brakhage, A.A., Kaveri, S.V., Romani, L., Latgé, J.-P., 2009. Surface hydrophobin prevents immune recognition of airborne fungal spores. *Nature* 460, 1117–1121. <https://doi.org/10.1038/nature08264>
- Armenante, A., Longobardi, S., Rea, I., De Stefano, L., Giocondo, M., Silipo, A., Molinaro, A., Giardina, P., 2010. The *Pleurotus ostreatus* hydrophobin Vmh2 and its interaction with glucans. *Glycobiology* 20, 594–602. <https://doi.org/10.1093/glycob/cwq009>
- Artini, M., Cicatiello, P., Ricciardelli, A., Papa, R., Selan, L., Dardano, P., Tilotta, M., Vrenna, G., Tutino, M.L., Giardina, P., Parrilli, E., 2017. Hydrophobin coating prevents *Staphylococcus epidermidis* biofilm formation on different surfaces. *Biofouling* 33, 601–611. <https://doi.org/10.1080/08927014.2017.1338690>
- Bayry, J., Beaussart, A., Dufrière, Y.F., Sharma, M., Bansal, K., Knemeyer, O., Aimanianda, V., Brakhage, A.A., Kaveri, S.V., Kwon-Chung, K.J., Latgé, J.-P., Beauvais, A., 2014. Surface structure characterization of *Aspergillus fumigatus* conidia mutated in the melanin synthesis pathway and their human cellular immune response. *Infect. Immun.* 82, 3141–3153. <https://doi.org/10.1128/IAI.01726-14>
- Beauvais, A., Bozza, S., Knemeyer, O., Formosa, Cécile, Formosa, Céline, Balloy, V., Henry, C., Roberson, R.W., Dague, E., Chignard, M., Brakhage, A.A., Romani, L., Latgé, J.-P., 2013. Deletion of the α -(1,3)-glucan synthase genes induces a restructuring of the conidial cell wall responsible for the avirulence of *Aspergillus fumigatus*. *PLoS Pathog.* 9, e1003716. <https://doi.org/10.1371/journal.ppat.1003716>
- Beauvais, A., Monod, M., Debeaupuis, J.P., Diaquin, M., Kobayashi, H., Latgé, J.P., 1997. Biochemical and antigenic characterization of a new dipeptidyl-peptidase isolated from *Aspergillus fumigatus*. *J. Biol. Chem.* 272, 6238–6244.
- Brunger, A.T., Adams, P.D., Clore, G.M., DeLano, W.L., Gros, P., Grosse-Kunstleve, R.W., Jiang, J.S., Kuszewski, J., Nilges, M., Pannu, N.S., Read, R.J., Rice, L.M., Simonson, T., Warren, G.L., 1998. Crystallography & NMR system: A new software suite for macromolecular structure determination. *Acta Crystallogr Biol Crystallogr* 54, 905–21.
- Bruns, S., Knemeyer, O., Hasenberg, M., Aimanianda, V., Nietzsche, S., Thywissen, A., Jeron, A., Latgé, J.-P., Brakhage, A.A., Gunzer, M., 2010. Production of extracellular traps against *Aspergillus fumigatus* in vitro and in infected lung tissue is dependent on invading neutrophils and influenced by hydrophobin RodA. *PLoS Pathog.* 6, e1000873. <https://doi.org/10.1371/journal.ppat.1000873>
- Carrion, S. de J., Leal, S.M., Ghannoum, M.A., Aimanianda, V., Latgé, J.-P., Pearlman, E., 2013. The RodA hydrophobin on *Aspergillus fumigatus* spores masks dectin-1- and dectin-2-dependent responses and enhances fungal survival in vivo. *J. Immunol. Baltim. Md 1950* 191, 2581–2588. <https://doi.org/10.4049/jimmunol.1300748>
- Chen, V.B., Arendall, W.B., Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S., Richardson, D.C., 2010. MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr Biol Crystallogr* 66, 12–21. <https://doi.org/10.1107/S0907444909042073>
- Cicatiello, P., Dardano, P., Pirozzi, M., Gravagnuolo, A.M., De Stefano, L., Giardina, P., 2017. Self-assembly of two hydrophobins from marine fungi affected by interaction with surfaces. *Biotechnol. Bioeng.* 114, 2173–2186. <https://doi.org/10.1002/bit.26344>
- Cox, J., Mann, M., 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* 26, 1367–1372. <https://doi.org/10.1038/nbt.1511>

- Cox, J., Neuhauser, N., Michalski, A., Scheltema, R.A., Olsen, J.V., Mann, M., 2011. Andromeda: a peptide search engine integrated into the MaxQuant environment. *J. Proteome Res.* 10, 1794–1805. <https://doi.org/10.1021/pr101065j>
- da Silva Ferreira, M.E., Kress, M.R.V.Z., Savoldi, M., Goldman, M.H.S., Härtl, A., Heinekamp, T., Brakhage, A.A., Goldman, G.H., 2006. The akuB(KU80) mutant deficient for nonhomologous end joining is a powerful tool for analyzing pathogenicity in *Aspergillus fumigatus*. *Eukaryot. Cell* 5, 207–211. <https://doi.org/10.1128/EC.5.1.207-211.2006>
- Dagenais, T.R.T., Giles, S.S., Aïmanianda, V., Latgé, J.-P., Hull, C.M., Keller, N.P., 2010. *Aspergillus fumigatus* LaeA-mediated phagocytosis is associated with a decreased hydrophobin layer. *Infect. Immun.* 78, 823–829. <https://doi.org/10.1128/IAI.00980-09>
- Dague, E., Alsteens, D., Latgé, J.-P., Dufrêne, Y.F., 2008a. High-resolution cell surface dynamics of germinating *Aspergillus fumigatus* conidia. *Biophys. J.* 94, 656–660. <https://doi.org/10.1529/biophysj.107.116491>
- Dague, E., Alsteens, D., Latgé, J.-P., Verbelen, C., Raze, D., Baulard, A.R., Dufrêne, Y.F., 2007. Chemical force microscopy of single live cells. *Nano Lett.* 7, 3026–3030. <https://doi.org/10.1021/nl071476k>
- Dague, E., Delcorte, A., Latgé, J.-P., Dufrêne, Y.F., 2008b. Combined use of atomic force microscopy, X-ray photoelectron spectroscopy, and secondary ion mass spectrometry for cell surface analysis. *Langmuir ACS J. Surf. Colloids* 24, 2955–2959. <https://doi.org/10.1021/la703741y>
- Gandier, J.-A., Langelaan, D.N., Won, A., O'Donnell, K., Grondin, J.L., Spencer, H.L., Wong, P., Tillier, E., Yip, C., Smith, S.P., Master, E.R., 2017. Characterization of a Basidiomycota hydrophobin reveals the structural basis for a high-similarity Class I subdivision. *Sci. Rep.* 7, 45863. <https://doi.org/10.1038/srep45863>
- Gastebois, A., Aïmanianda, V., Bachellier-Bassi, S., Nesseir, A., Firon, A., Beauvais, A., Schmitt, C., England, P., Beau, R., Prévost, M.-C., d'Enfert, C., Latgé, J.-P., Mouyna, I., 2013. SUN proteins belong to a novel family of β -(1,3)-glucan-modifying enzymes involved in fungal morphogenesis. *J. Biol. Chem.* 288, 13387–13396. <https://doi.org/10.1074/jbc.M112.440172>
- Gebbink, M.F.B.G., Claessen, D., Bouma, B., Dijkhuizen, L., Wösten, H.A.B., 2005. Amyloids — a functional coat for microorganisms. *Nat. Rev. Microbiol.* 3, 333–341. <https://doi.org/10.1038/nrmicro1127>
- Grünbacher, A., Throm, T., Seidel, C., Gutt, B., Röhrig, J., Strunk, T., Vincze, P., Walheim, S., Schimmel, T., Wenzel, W., Fischer, R., 2014. Six hydrophobins are involved in hydrophobin rodlet formation in *Aspergillus nidulans* and contribute to hydrophobicity of the spore surface. *PloS One* 9, e94546. <https://doi.org/10.1371/journal.pone.0094546>
- Halle, A., Hornung, V., Petzold, G.C., Stewart, C.R., Monks, B.G., Reinheckel, T., Fitzgerald, K.A., Latz, E., Moore, K.J., Golenbock, D.T., 2008. The NALP3 inflammasome is involved in the innate immune response to amyloid-beta. *Nat. Immunol.* 9, 857–865. <https://doi.org/10.1038/ni.1636>
- Hartmann, T., Dümig, M., Jaber, B.M., Szewczyk, E., Olbermann, P., Morschhäuser, J., Krappmann, S., 2010. Validation of a self-excising marker in the human pathogen *Aspergillus fumigatus* by employing the beta-rec/six site-specific recombination system. *Appl. Environ. Microbiol.* 76, 6313–6317. <https://doi.org/10.1128/AEM.00882-10>
- Hektor, H.J., Scholtmeijer, K., 2005. Hydrophobins: proteins with potential. *Curr. Opin. Biotechnol.* 16, 434–439. <https://doi.org/10.1016/j.copbio.2005.05.004>

- Hooft, R.W., Vriend, G., Sander, C., Abola, E.E., 1996. Errors in protein structures. *Nature* 381, 272. <https://doi.org/10.1038/381272a0>
- Jensen, B.G., Andersen, M.R., Pedersen, M.H., Frisvad, J.C., Søndergaard, I., 2010. Hydrophobins from *Aspergillus* species cannot be clearly divided into two classes. *BMC Res. Notes* 3, 344. <https://doi.org/10.1186/1756-0500-3-344>
- Kay, L.E., Xu, G., Singer, A.U., Muhandiram, D.R., Forman-Kay, J.D., 1993. A gradient-enhanced HCCH-TOCSY experiment for recording side-chain ^1H and ^{13}C correlations in H_2O samples of proteins. *J Magn Reson Ser. B* 101, 333–337.
- Kershaw, M.J., Talbot, N.J., 1998. Hydrophobins and repellents: proteins with fundamental roles in fungal morphogenesis. *Fungal Genet. Biol. FG B* 23, 18–33. <https://doi.org/10.1006/fgbi.1997.1022>
- Kershaw, M.J., Wakley, G., Talbot, N.J., 1998. Complementation of the *mpg1* mutant phenotype in *Magnaporthe grisea* reveals functional relationships between fungal hydrophobins. *EMBO J.* 17, 3838–3849. <https://doi.org/10.1093/emboj/17.14.3838>
- Kwan, A.H.Y., Winefield, R.D., Sunde, M., Matthews, J.M., Haverkamp, R.G., Templeton, M.D., Mackay, J.P., 2006. Structural basis for rodlet assembly in fungal hydrophobins. *Proc. Natl. Acad. Sci. U. S. A.* 103, 3621–3626. <https://doi.org/10.1073/pnas.0505704103>
- Lambou, K., Lamarre, C., Beau, R., Dufour, N., Latgé, J.-P., 2010. Functional analysis of the superoxide dismutase family in *Aspergillus fumigatus*. *Mol. Microbiol.* 75, 910–923. <https://doi.org/10.1111/j.1365-2958.2009.07024.x>
- Laskowski, R.A., MacArthur, M.W., Moss, D.S., Thornton, J.M., 1993. PROCHECK: a program to check the stereochemical quality of protein structures. *J Appl Crystallogr* 26, 283–291.
- Latgé, J.P., 1999. *Aspergillus fumigatus* and aspergillosis. *Clin. Microbiol. Rev.* 12, 310–350.
- Latgé, J.-P., Beauvais, A., Chamilos, G., 2017. The Cell Wall of the Human Fungal Pathogen *Aspergillus fumigatus*: Biosynthesis, Organization, Immune Response, and Virulence. *Annu. Rev. Microbiol.* 71, 99–116. <https://doi.org/10.1146/annurev-micro-030117-020406>
- littlejohn, K.A., Hooley, P., Cox, P. w., 2012. Bioinformatics predicts diverse *Aspergillus* hydrophobins with novel properties. *Food Hydrocoll.* 27, 503–516. <https://doi.org/10.1016/j.foodhyd.2011.08.018>
- Longobardi, S., Picone, D., Ercole, C., Spadaccini, R., De Stefano, L., Rea, I., Giardina, P., 2012. Environmental conditions modulate the switch among different states of the hydrophobin *Vmh2* from *Pleurotus ostreatus*. *Biomacromolecules* 13, 743–750. <https://doi.org/10.1021/bm201663f>
- Macindoe, I., Kwan, A.H., Ren, Q., Morris, V.K., Yang, W., Mackay, J.P., Sunde, M., 2012. Self-assembly of functional, amphipathic amyloid monolayers by the fungal hydrophobin EAS. *Proc. Natl. Acad. Sci. U. S. A.* 109, E804–811. <https://doi.org/10.1073/pnas.1114052109>
- Melin, P., Schnürer, J., Wagner, E.G.H., 2003. Characterization of *phiA*, a gene essential for phialide development in *Aspergillus nidulans*. *Fungal Genet. Biol. FG B* 40, 234–241.
- Monod, M., Capoccia, S., Léchenne, B., Zaugg, C., Holdom, M., Jousson, O., 2002. Secreted proteases from pathogenic fungi. *Int. J. Med. Microbiol. IJMM* 292, 405–419. <https://doi.org/10.1078/1438-4221-00223>
- Morris, V.K., Linser, R., Wilde, K.L., Duff, A.P., Sunde, M., Kwan, A.H., 2012. Solid-state NMR spectroscopy of functional amyloid from a fungal hydrophobin: a well-ordered β -sheet core amidst structural heterogeneity. *Angew. Chem. Int. Ed Engl.* 51, 12621–12625. <https://doi.org/10.1002/anie.201205625>

- Morris, V.K., Ren, Q., Macindoe, I., Kwan, A.H., Byrne, N., Sunde, M., 2011. Recruitment of class I hydrophobins to the air:water interface initiates a multi-step process of functional amyloid formation. *J. Biol. Chem.* 286, 15955–15963. <https://doi.org/10.1074/jbc.M110.214197>
- Muhandiram, D.R., Kay, L.E., 1994. Gradient-enhanced triple-resonance three-dimensional NMR experiments with improved sensitivity. *J Magn Reson Ser. B* 103, 203–216.
- Niu, B., Li, B., Wang, H., Guo, R., Xu, H., Qiao, M., Li, W., 2017. Investigation of the relationship between the rodlet formation and Cys3-Cys4 loop of the HGFI hydrophobin. *Colloids Surf. B Biointerfaces* 150, 344–351. <https://doi.org/10.1016/j.colsurfb.2016.10.048>
- Paris, S., Debeaupuis, J.-P., Crameri, R., Carey, M., Charlès, F., Prévost, M.C., Schmitt, C., Philippe, B., Latgé, J.P., 2003. Conidial hydrophobins of *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* 69, 1581–1588.
- Pedersen, M.H., Borodina, I., Moresco, J.L., Svendsen, W.E., Frisvad, J.C., Søndergaard, I., 2011. High-yield production of hydrophobins RodA and RodB from *Aspergillus fumigatus* in *Pichia pastoris*. *Appl. Microbiol. Biotechnol.* 90, 1923–1932. <https://doi.org/10.1007/s00253-011-3235-1>
- Petersen, T.N., Brunak, S., von Heijne, G., Nielsen, H., 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods* 8, 785–786. <https://doi.org/10.1038/nmeth.1701>
- Pham, C.L.L., Rey, A., Lo, V., Soulès, M., Ren, Q., Meisl, G., Knowles, T.P.J., Kwan, A.H., Sunde, M., 2016. Self-assembly of MPG1, a hydrophobin protein from the rice blast fungus that forms functional amyloid coatings, occurs by a surface-driven mechanism. *Sci. Rep.* 6, 25288. <https://doi.org/10.1038/srep25288>
- Pille, A., Kwan, A.H., Cheung, I., Hampsey, M., Aimanianda, V., Delepierre, M., Latgé, J.-P., Sunde, M., Guijarro, J.I., 2015. (1)H, (13)C and (15)N resonance assignments of the RodA hydrophobin from the opportunistic pathogen *Aspergillus fumigatus*. *Biomol. NMR Assign.* 9, 113–118. <https://doi.org/10.1007/s12104-014-9555-1>
- Piscitelli, A., Cicatiello, P., Gravagnuolo, A.M., Sorrentino, I., Pezzella, C., Giardina, P., 2017. Applications of Functional Amyloids from Fungi: Surface Modification by Class I Hydrophobins. *Biomolecules* 7. <https://doi.org/10.3390/biom7030045>
- Reichard, U., Monod, M., Odds, F., Rüchel, R., 1997. Virulence of an aspergillopepsin-deficient mutant of *Aspergillus fumigatus* and evidence for another aspartic proteinase linked to the fungal cell wall. *J. Med. Vet. Mycol. Bi-Mon. Publ. Int. Soc. Hum. Anim. Mycol.* 35, 189–196.
- Ren, Q., Kwan, A.H., Sunde, M., 2013. Two forms and two faces, multiple states and multiple uses: Properties and applications of the self-assembling fungal hydrophobins. *Biopolymers* 100, 601–612. <https://doi.org/10.1002/bip.22259>
- Ribitsch, D., Herrero Acero, E., Przylucka, A., Zitzenbacher, S., Marold, A., Gamerith, C., Tscheließnig, R., Jungbauer, A., Rennhofer, H., Lichtenegger, H., Amenitsch, H., Bonazza, K., Kubicek, C.P., Druzhinina, I.S., Guebitz, G.M., 2015. Enhanced cutinase-catalyzed hydrolysis of polyethylene terephthalate by covalent fusion to hydrophobins. *Appl. Environ. Microbiol.* 81, 3586–3592. <https://doi.org/10.1128/AEM.04111-14>
- Rieping, W., Habeck, M., Bardiaux, B., Bernard, A., Malliavin, T.E., Nilges, M., 2007. ARIA2: automated NOE assignment and data integration in NMR structure calculation. *Bioinformatics* 23, 381–382.
- Sarfati, J., Monod, M., Recco, P., Sulahian, A., Pinel, C., Candolfi, E., Fontaine, T., Debeaupuis, J.-P., Tabouret, M., Latgé, J.-P., 2006. Recombinant antigens as

- diagnostic markers for aspergillosis. *Diagn. Microbiol. Infect. Dis.* 55, 279–291. <https://doi.org/10.1016/j.diagmicrobio.2006.02.002>
- Scholtmeijer, K., de Vocht, M.L., Rink, R., Robillard, G.T., Wösten, H.A.B., 2009. Assembly of the fungal SC3 hydrophobin into functional amyloid fibrils depends on its concentration and is promoted by cell wall polysaccharides. *J. Biol. Chem.* 284, 26309–26314. <https://doi.org/10.1074/jbc.M109.005553>
- Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., Selbach, M., 2011. Global quantification of mammalian gene expression control. *Nature* 473, 337–342. <https://doi.org/10.1038/nature10098>
- Sharma, D., Rajarathnam, K., 2000. ¹³C NMR chemical shifts can predict disulfide bond formation. *J. Biomol. NMR* 18, 165–171.
- Shen, Y., Delaglio, F., Cornilescu, G., Bax, A., 2009. TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *J. Biomol. NMR* 44, 213–23. <https://doi.org/10.1007/s10858-009-9333-z>
- Sippl, M.J., 1993. Recognition of errors in three-dimensional structures of proteins. *Proteins* 17, 355–62. <https://doi.org/10.1002/prot.340170404>
- Sriranganadane, D., Waridel, P., Salamin, K., Reichard, U., Grouzmann, E., Neuhaus, J.-M., Quadroni, M., Monod, M., 2010. Aspergillus protein degradation pathways with different secreted protease sets at neutral and acidic pH. *J. Proteome Res.* 9, 3511–3519. <https://doi.org/10.1021/pr901202z>
- Sunde, M., Kwan, A.H.Y., Templeton, M.D., Beever, R.E., Mackay, J.P., 2008. Structural analysis of hydrophobins. *Micron Oxf. Engl.* 1993 39, 773–784. <https://doi.org/10.1016/j.micron.2007.08.003>
- Takahashi, T., Maeda, H., Yoneda, S., Ohtaki, S., Yamagata, Y., Hasegawa, F., Gomi, K., Nakajima, T., Abe, K., 2005. The fungal hydrophobin RolA recruits polyesterase and laterally moves on hydrophobic surfaces. *Mol. Microbiol.* 57, 1780–1796. <https://doi.org/10.1111/j.1365-2958.2005.04803.x>
- Talbot, N.J., Ebole, D.J., Hamer, J.E., 1993. Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant Cell* 5, 1575–1590. <https://doi.org/10.1105/tpc.5.11.1575>
- Talbot, N.J., Kershaw, M.J., Wakley, G.E., De Vries, O.M.H., Wessels, J.G.H., Hamer, J.E., 1996. MPG1 Encodes a Fungal Hydrophobin Involved in Surface Interactions during Infection-Related Development of *Magnaporthe grisea*. *Plant Cell* 8, 985–999. <https://doi.org/10.1105/tpc.8.6.985>
- Tanaka, T., Nakayama, M., Takahashi, T., Nanatani, K., Yamagata, Y., Abe, K., 2017. Analysis of the ionic interaction between the hydrophobin RodA and two cutinases of *Aspergillus nidulans* obtained via an *Aspergillus oryzae* expression system. *Appl. Microbiol. Biotechnol.* 101, 2343–2356. <https://doi.org/10.1007/s00253-016-7979-5>
- Thau, N., Monod, M., Crestani, B., Rolland, C., Tronchin, G., Latgé, J.P., Paris, S., 1994. rodletless mutants of *Aspergillus fumigatus*. *Infect. Immun.* 62, 4380–4388.
- Tsolis, A.C., Papandreou, N.C., Iconomidou, V.A., Hamodrakas, S.J., 2013. A consensus method for the prediction of “aggregation-prone” peptides in globular proteins. *PLoS One* 8, e54175. <https://doi.org/10.1371/journal.pone.0054175>
- Tükel, C., Wilson, R.P., Nishimori, J.H., Pezeshki, M., Chromy, B.A., Bäumlner, A.J., 2009. Responses to amyloids of microbial and host origin are mediated through toll-like receptor 2. *Cell Host Microbe* 6, 45–53. <https://doi.org/10.1016/j.chom.2009.05.020>
- Tyanova, S., Temu, T., Cox, J., 2016. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat. Protoc.* 11, 2301–2319. <https://doi.org/10.1038/nprot.2016.136>

- Valsecchi, I., Dupres, V., Stephen-Victor, E., Guijarro, J.I., Gibbons, J., Beau, R., Bayry, J., Coppee, J.-Y., Lafont, F., Latgé, J.-P., Beauvais, A., 2017. Role of Hydrophobins in *Aspergillus fumigatus*. *J. Fungi Basel Switz.* 4. <https://doi.org/10.3390/jof4010002>
- Wang, X., Permentier, H.P., Rink, R., Kruijtzter, J. a. W., Liskamp, R.M.J., Wösten, H. a. B., Poolman, B., Robillard, G.T., 2004. Probing the self-assembly and the accompanying structural changes of hydrophobin SC3 on a hydrophobic surface by mass spectrometry. *Biophys. J.* 87, 1919–1928. <https://doi.org/10.1529/biophysj.104.041616>
- Whiteford, J.R., Spanu, P.D., 2002. Hydrophobins and the interactions between fungi and plants. *Mol. Plant Pathol.* 3, 391–400. <https://doi.org/10.1046/j.1364-3703.2002.00129.x>
- Zhang, S., Xia, Y.X., Kim, B., Keyhani, N.O., 2011. Two hydrophobins are involved in fungal spore coat rodlet layer assembly and each play distinct roles in surface interactions, development and pathogenesis in the entomopathogenic fungus, *Beauveria bassiana*. *Mol. Microbiol.* 80, 811–826. <https://doi.org/10.1111/j.1365-2958.2011.07613.x>
- Zhao, Z.-X., Wang, H.-C., Qin, X., Wang, X.-S., Qiao, M.-Q., Anzai, J.-I., Chen, Q., 2009. Self-assembled film of hydrophobins on gold surfaces and its application to electrochemical biosensing. *Colloids Surf. B Biointerfaces* 71, 102–106. <https://doi.org/10.1016/j.colsurfb.2009.01.011>
- Zykwinska, A., Guillemette, T., Bouchara, J.-P., Cuenot, S., 2014a. Spontaneous self-assembly of SC3 hydrophobins into nanorods in aqueous solution. *Biochim. Biophys. Acta* 1844, 1231–1237. <https://doi.org/10.1016/j.bbapap.2014.04.003>
- Zykwinska, A., Pihet, M., Radji, S., Bouchara, J.-P., Cuenot, S., 2014b. Self-assembly of proteins into a three-dimensional multilayer system: investigation of the surface of the human fungal pathogen *Aspergillus fumigatus*. *Biochim. Biophys. Acta* 1844, 1137–1144. <https://doi.org/10.1016/j.bbapap.2014.03.001>

Table 1

The spreadsheet "Pres115146_AbsKu80_30d2h" gives the list of proteins with a signal peptide extracted with formic acid from 30 days old conidia, which are specific of the double mutant conidia I115S/I146G (absent from the parental Ku80 strain); the spreadsheet "Pres115146_AbsKu80_10d2h" gives the list of proteins with a signal peptide extracted with formic acid from 10 days old conidia, which are specific of the double mutant conidia I115S/I146G (and absent from the Ku80 parental strain); the spread sheet "Common_prot-30d2h" ranks (from the most abundant to the least abundant) the 31 proteins with signal peptides which are common in the formic acid extracts from 10 days and 30 days old conidia. Over 1000 proteins were identified in the formic acid extracts from each strain and were deposited in the PRIDE data base (<https://www.ebi.ac.uk/pride/>).

Figures legends

Figure 1. Structure (A), internal dynamics (B) and hydrophobicity (C) of monomeric RodA (best energy conformer). **A:** Strands are shown in blue, helices in light blue and disulfide bridges in marine blue. The inter cysteine regions on the front (black) and back (grey) as well as the N-terminal region are indicated. Cysteines are numbered by their order in the sequence throughout this paper. **B:** Backbone amide ^1H - ^{15}N heteronuclear $n\text{Oe}$ values are shown on the cartoon representation of the structure: highly flexible ($n\text{Oe} < 0.45$), flexible ($0.45 \leq n\text{Oe} \leq 0.65$), rigid ($n\text{Oe} > 0.65$) and missing-value backbone residues are shown in red, orange, blue and grey, respectively. The structure is shown in a different orientation on the right. **C:** The hydropathy profile calculated with the Eisenberg hydrophobicity scale (<http://web.expasy.org/protscale>) is color coded on the surface and cartoon representations of the structure: blue= hydrophilic ($\Phi \leq -0.3$), orange ($-0.3 < \Phi < 0.3$), red=hydrophobic ($\Phi \geq 0.3$). The N-terminal disordered region (residues 18-38) is not displayed in **B** and **C**.

Figure 2. The effect of mutation(s) on the kinetics of RodA rodlet self-assembly assessed by Thioflavin T fluorescence at 50 °C. Amyloid assembly profiles for **(A)** RodA WT; single point mutants **(B)** RodA I114G, **(C)** RodA I115G, **(D)** RodA I145G and **(E)** RodA I146G; **(F)** double point mutant RodA I115G-I146G and **(G)** C7C8NC2 chimera. **(H)** Lag time and **(I)** time taken to reach half maximum ThT fluorescence. (See the material and methods section for calculation of lag time and time to reach half maximum).

Figure 3. Assembly of RodA WT and mutant proteins on highly oriented pyrolytic graphite. Atomic force micrographs showing the surface morphology of rodlets formed by **(A)** RodA WT, **(B)** RodA I114G, **(C)** RodA I115G, **(D)** RodA I145G and **(E)** RodA I146G after incubation at room temperature on graphite surfaces. Surface layers containing rodlets formed by **(F)** RodA WT, **(G)** double mutant RodA I115G/I146G and **(H)** C7C8NC2 chimera after incubation at 50°C for 2 hours. (See the Material and Methods section for the AFM experimental conditions).

Figure 4. Transmission electron micrographs of fibrillar material formed by **(A)** PIIGIPIQDL, a peptide derived from the C4-C5 region of RodA and **(B)** SLIGL, a peptide with predicted amyloidogenic sequence from the C7-C8 region.

Figure 5. Cysteine mutations impair the presence of rodlets at the conidial surface. AFM deflection images showing the structure of conidial surface imaged in liquid conditions for: **(A)** the parental, **(B)** one cysteine (C127S) and **(C)** four cysteine (C64S/C65S/C133S/C134S) mutants.

Figure 6. Delay in rodlets appearance on the surface of conidia imaged in liquid conditions. AFM deflection images of the rodlets layer for (A-C) the parental, (D-F) I146G and (G-I) I115S/I146G mutants after 3 (A, D, G), 10 (B, E, H) and 30 (C, F, I) days of growth.

Figure 7. The surface of the RodA I115S/I146G double mutant is only partially covered by rodlets. AFM deflections images (at different magnification) representative of the conidium surface of the RodA I115S/I146G double mutant after 30 days of culture with different magnifications.

Figure 8. Conidia of single and quadruple cysteine mutants and the double RodA I115S/I146G mutant activate the human immune response.

A-B: Induction of the dendritic cells (DC) maturation. **C-D:** Production of cytokines by DC. **E-F:** Polarization of T cell response. As other single point mutants did not alter either DC phenotype or cytokines, they were not tested for T cell response. Note that (i) the cysteine mutants also polarize the Treg response and that (ii) the I146G mutant slightly activates the DC response.

Figure 9. Lack of DC activation by the reduced (A,B) or mutated RodA protein (C,D).

A-B: Unfolded rRodA, with or without reduced and blocked Cys residues with N-ethyl maleimide (NEM) or iodoacetamide (IA) do not activate dendritic cells (A) and do not induce cytokine production (B). **C-D:** RodA extracted from conidia of mutants L145S, I146G, I115G/I146S (C) or lacking the N terminal region (RodA-Nterm) (D), are immunologically inert as parental RodA. LPS is used as a control.

Figure 10. The cell wall Pep1 and Pep2 aspartic proteases (used as recombinant proteins) specifically degraded the RodA protein (see Material and Methods for the experimental conditions).

Supplementary Tables

Table S1: Statistics for the ensemble of 10 structures calculated for RodA (19-159).

Table S2. Width and height of the rodlets formed by wild type and mutated rRodA (measured from cross-sectional analysis at multiple sites across the AFM images).

Table S3. Primers used for the construction of point mutated RodA *A. fumigatus* mutants and aspartic protease Δ *ctsD*, Δ *opsB*, Δ *pep1* and Δ *pep2* mutants.

Supplementary Figures

Figure S1. A, ^1H - ^{15}N HSQC of recombinant RodA (blue) and B comparison with the spectrum of the protein HF-extracted from spores (magenta). Spectra were recorded at 25°C in 50 mM CD_3COONa pH 4.3 10% D_2O and referenced to DSS (sodium salt of 4,4-dimethyl-4-silapentane-1-sulfonic acid). In **A**, the assignments of all the backbone amide protons are displayed. For the sake of simplicity, the side chain amide and amino group signals are not labeled. In **B**, only the signals of the amide groups of the 8 Cys residues are labeled and the C-terminal NH signal (159L) is indicated by an arrow. Although the HF-extracted protein was a mixture of folded and unfolded forms that we could not resolve by reverse phase chromatography using analytical c18 columns and showed many broad peaks suggesting heterogeneity in the sample, the comparison of the NMR spectra clearly indicate that both proteins show the same structure. Based on the assignments of rRodA, we assigned the backbone resonances of the extracted protein between residues 23 and 159.

Figure S2. Structure ensemble of RodA. The ensemble consists of the 10 best energy conformers. β -sheets, α -helices and disulfide bonds are colored in magenta, cyan and blue marine, respectively.

Figure S3. Backbone amide heteronuclear ^1H - ^{15}N nOe parameter as a function of residue. Three root mean-square deviations (rmsd) of noise were used to estimate the error in peak intensities. The secondary structure (red) and cysteine positions (grey) are shown. Cylinders represent α or 3_{10} -helices and arrows β -strands. Missing values are due to Pro residues or signal overlap for Q111.

Figure S4. Construction of the *A. fumigatus* mutant strain lacking the N-terminal region (RodA-Nterm). **A.** Representation of the DNA fragments used to construct the RodA-Nterm cassette. **B.** Diagram showing the *Bam*HI digestion site and the DIG-labeled probe location. **C.** Southern blot of the parental and RodA-Nterm strains. **D.** AFM image showing the normal rodlet morphology on the conidial surface of the mutant.

Figure S5. Construction of *A. fumigatus* point mutated RodA strains. **A.** The cassette used to knock out *Roda* was constructed to replace the *Roda* ORF by the point mutated *RODA* linked to a hygromycin resistance β -recombinase cassette (*HPH*^R- β -rec) through double crossing-over of the upstream and downstream borders (blue-shaded boxes). The *Bst*Z171 restriction sites are positioned in the diagram. **B.** Southern blot analysis of gDNA of parental and mutated RodA strains digested with *Bst*Z171 and probed with a DIG-labeled probe of the *RODA* sequence.

Figure S6. RodA is not present on the conidial surface of the RodA cysteine-mutated strains (C127S or C64S/C65S/C133S/C134S). Conidial proteins extracted with HF from parental and RodA point mutated in one (C127S) or four cysteines (C64S/C65S/C133S/C134S) were separated on SDS-PAGE and analyzed by western blot with antibodies against RodA.

Figure S7. Conidial proteins extracted with formic acid from 30-days old from parental and RodA-mutated conidia. A. Total proteins stained with colloidal Coomassie blue. Note the enrichment in proteins extracted by HF in the mutants without rodlet ($\Delta rodA$ or cysteine mutants). **B.** Western blot using an anti-RodA antibody. Note that only the lower Mr band is seen in the mutant (RodA-Nterm) that lacks the N-terminal residues 19-39.

Figure S8. Length of the rodlets of the I146G and I115S/I146G RodA mutants and parental strain. The length of the parental and single mutants are similar and significantly ($p < 0.01$) shorter than the I146G and I115S/I146G mutants. *: Significant differences.

Figure S9. Labelling of the conidial surface of the I115S/I146S mutant with ConA-FITC or WGA-FITC. Note the absence of label in parental conidia.

Figure S10. Phenotypes of point mutated RodA conidia. RodA C127S, RodA C64S/C65S/C133S/C134S and $\Delta rodA$ show typical black conidia observed in the absence of RodA at the conidial surface. Similarly, the RodA I115S/I146S strain, which has a small proportion of the surface covered by rodlets, shows black conidia. The RodA I146S strain also displays darker conidia than the parental strain.

Figure S11. Degradation of the rodlet protein induced by the addition of a culture filtrate (BCAP) of germinating conidia. Degradation of the RodA protein doublet is inhibited by the addition of pepstatin or heat inactivation. Contrastingly, inhibitors of the major *A. fumigatus* secreted serine proteases (PMSF, TLCK) do not block the degradation.

Figure S12. Construction of the *A. fumigatus* $\Delta ctsD$ strain. A. The DNA construct used to knock out *CTSD* was used to replace the ORF of the *CTSD* sequence by the hygromycin resistance β -recombinase cassette (HPH^R β -rec), through double crossing-over of the upstream and downstream borders. **B.** Southern blot analysis. The gDNA of parental and $\Delta ctsD$ strains were digested with *XhoI* (restriction site positioned in diagram A) and probed with a DIG-labeled probe containing the upstream border as shown in the diagram.

Figure S13. Construction of the *A. fumigatus* $\Delta opsB$ strain. A. *OPSB* was replaced by the hygromycin resistance β -recombinase cassette (HPH^R β -rec) through double crossing-over of the upstream and downstream borders. **B.** Southern blot analysis. The gDNA of parental and $\Delta opsB$ strains were digested with *EcoRI* (restriction site positioned in diagram A) and probed with a DIG-labeled probe of the upstream border.

Figure S14. Construction of the *A. fumigatus* $\Delta pep1$ strain. A. *PEP1* was replaced by the hygromycin resistance β -recombinase cassette (HPH^R β -rec) through double crossing-over of the upstream and downstream borders. **B.** Southern blot analysis. The gDNA of parental and $\Delta pep1$ strains were digested with *ClaI* (restriction site positioned in in diagram A) and probed with a DIG-labeled probe containing the downstream border as shown in the diagram.

Figure S15. Construction of the *A. fumigatus* $\Delta pep2$ strain. **A.** *PEP2* was replaced by the hygromycin resistance β -recombinase cassette (HPH^R β -rec) through double crossing-over of the upstream and downstream borders. **B.** Southern blot analysis. The gDNA of the parental and $\Delta pep2$ strains were digested with *Bam*HI (restriction site positioned in diagram A) probed with a DIG-labeled probe containing the upstream border as shown in the diagram.

ACCEPTED MANUSCRIPT

Proteins with signal peptide which are quantified in 115146 and not in Ku80 at 30d and 2h of extraction (Ranked by average iBAQ intensity in 115146)	Statistics (rank from the highest mean of iBAQ values to the lowest to rank from the most abundant protein to the least)	Number of identified peptides specific to each protein in each sample	iBAQ (intensity-based absolute quantification) values in each sample
<p>Fasta.headers</p> <p>UPs PUSM nret rneo isr oiq pe i Btqu rnt n Lue oing Awenw t1 _ Si . ce G . 1 p Tts ei ea 5 r _ he . g nc 1 o A ql h ec 4 t f s u e t . e 6 e 2 i e n . l s a i 9 g n g . D s n n 3 n c t k</p>	<p>MM aa nn Bi Aq Qu ep 3p 0e dp 2ti</p>	<p>UUUUUU nnnnnn iiiii qqqqqq uuuuuu eeeee pppppp pppppp tittitit d d d d d e e e e e s. s. s. s. s.</p>	<p>iiiii BBBBA AAQA QQQQQ 33333 00000 ddddd _ _ _ _ _ 22222 hhhhh _ _ _ _ _ 111D D D 111e e e</p>

i	d	_	a	e	h	D	h	d	3	3	3	3	3	3	5	5	5	l	l	l
o	a	i	l	.	.	a	1	e	0	0	0	0	0	0	.	.	.	t	t	t
n	b	d	c	.	.	.	1	s	d	d	d	d	d	d	1	1	1	a	a	a
s			p	o			5	3	-	-	-	-	-	-	4	4	4	K	K	K
e			p	e	v		1	0	2	2	2	2	2	2	6	6	6	U	U	U
nt			p	e	r		4	d	h	h	h	h	h	h	-	-	-	8	8	8
in			t	r			6	2	-	-	-	-	-	-	1	2	3	0	0	0
K			i	a				h	1	1	1	D	D	D				-	-	-
u			d	g				1	1	1	1	e	e	e				1	2	3
8			e	e				1	5	5	5	lt	lt	lt						
0				..				5	.	.	.	a	a	a						
at			(..				1	1	1	1	K	K	K						
1			S	c	o	r		4	4	4	4	U	U	U						
0			c	o	r	e		6	6	6	6	8	8	8						
d									-	-	-	0	0	0						
2									1	2	3	-	-	-						
h												1	2	3						

tr BOYBK6	A	B	5	A	0		2	1							1	1	1						
BOYBK6_AS	F	0	0	F	.	2	1	7							7	7	8						
PFC	U	Y	y	8	U	7	9	1	.	5					5	0	0						
Uncharacter	B	B	e	5	A	8	.	9	3	7					9	9	5						
ized protein	_	K	s	.	_	2	3	8	8	9	4	4	4	4	0	0	0	1	4	2	0	0	0

OS=Neosart	0	6	C	7			4	0											0	0	0
orya	8		A	G				0											0	0	0
fumigata	6		D	0				0											0	0	0
(strain	9		A	0																	
CEA10 / CBS	5		F	3																	
144.89 /	0		U	7																	
FGSC			B	0																	
A1163)			P																		
GN=AFUB_0			0																		
86950 PE=4			0																		
SV=1			0																		
			0																		
			8																		
			4																		
			5																		
			1																		
			5																		
			0																		
			8																		
			5																		
tr BOYDZ8			.																		
BOYDZ8_AS			C																		
PFC			A																		
Putative			D																		
uncharacter			A																		
ized protein			F	A																	
OS=Neosart	A		U	F																	
orya	F		B	U				1	2												
fumigata	U		P	A				7	.												
(strain	B		0	_				4	6										2	2	
CEA10 / CBS			0	6				2	6										6	0	5
144.89 /	0	B	0	G				1	5	6									5	0	7
FGSC	9	0	0	0	0			2	8	6									3	3	0
A1163)	7	Y	9	0	.	3		.	6	6									6	2	9
GN=AFUB_0	3	D	4	1	7	8	1	0	6	6									6	0	0
97339 PE=4	3	Z	n	6	8	3	.	0	5	.	6								0	0	0
SV=1	9	8	o	3	0	5	5	9	3	7	7	2	3	3	0	0	0	0	0	0	0
tr BOYDU9	A		5	A																	
BOYDU9_AS	F		0	F																	
PFC	U		8	U																	1
Putative	B		5	A				8											8	6	0
uncharacter	_	B	.	_				1	6										4	7	6
ized protein	0	0	C	6	0			0	4										9	9	3
OS=Neosart	9	Y	A	G	.	2		.	1										6	9	0
orya	6	D	D	0	8	1		8	8										0	4	0
fumigata	8	U	n	A	0	5	.	9	5	0									0	0	0
(strain	6	9	o	F	6	3	4	8	3	0	1	1	1	1	1	0	0	0	0	0	0

CEA10 / CBS	0	U	1																	
144.89 /		B	0																	
FGSC		P																		
A1163)		0																		
GN=AFUB_0		0																		
96860 PE=4		0																		
SV=1		0																		
		9																		
		4																		
		1																		
		4																		
		5																		
		0																		
		8																		
		5																		
		.																		
tr BOY1U6		C																		
BOY1U6_AS		A																		
PFC		D																		
Agmatinase,		A																		
putative		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U				7													
fumigata	U	P	A				9													
(strain	B	0	_				3													
CEA10 / CBS	_	0	5				2						6	8	8					
144.89 /	0	B	0	G			4	3					8	4	5					
FGSC	6	0	0	1	0		2	3					3	5	0					
A1163)	0	Y	5	3	. 2		.	3					5	8	3					
GN=AFUB_0	8	1	y	9	1	6	8	3	1				5	2	3					
60890 PE=3	9	U	e	4	8	7	.	9	7	3			0	0	0					
SV=1	0	6	s	8	0	8	4	1	9	3	6	6	6	6	0	0	0	0	0	0
tr BOY1D4		5																		
BOY1D4_AS		0																		
PFC		8																		
Disulfide		5	A																	
isomerase	A	.	F																	
(TigA),	F	C	U							1										
putative	U	A	A							4										
OS=Neosart	B	D	_							.										
orya	_	A	5				5	3					5	6	5					
fumigata	0	B	F	G			4	9	3				8	4	6					
(strain	5	0	U	1	0		0	6	3				2	4	2					
CEA10 / CBS	9	Y	B	2	. 5		.	5	3				6	9	0					
144.89 /	8	1	y	P	2	8	4	3	2	5	3		8	2	5					
FGSC	9	D	e	0	6	6	.	6	5	0	3	1	1	1				0	0	0
A1163)	0	4	s	0	0	7	6	8	3	0	3	4	5	4	0	0	0	0	0	0

cargo	—	S	.	—	7		3	5	3							1	8	8
receptor	0	2	C	4			9	6	3							0	0	0
(Erv25),	7		A	G				6	3							0	0	0
putative	0		D	1				.	3									
OS=Neosart	0		A	3				6	3									
orya	8		F	1				7	3									
fumigata	0		U	9														
(strain			B	0														
CEA10 / CBS			P															
144.89 /			0															
FGSC			0															
A1163)			0															
GN=AFUB_0			0															
70080 PE=3			6															
SV=1			8															
			2															
			8															
			5															
			0															
			8															
			5															
tr B0YCQ5			.															
B0YCQ5_AS			C															
PFC SUN			A															
domain			D															
protein			A															
(Uth1),			A															
putative			F	A														
OS=Neosart	A		U	F														
orya	F		B	U					5									
fumigata	U		P	A					.									
(strain	B		0	—														
CEA10 / CBS	—		0	7					2	3						2	1	2
144.89 /	0	B	0	G					4	1	3					5	4	3
FGSC	9	0	0	0	0				3	1	3					9	0	2
A1163)	1	Y	8	5	.	2			.	0	3					7	4	8
GN=AFUB_0	0	C	y	8	4	4	3	4	5	2	3					5	4	7
91030 PE=4	3	Q	e	5	5	9	.	1	0	0	3					0	0	0
SV=1	0	5	s	0	0	6	2	4	4	0	3	6	4	6	0	0	0	0
tr B0Y5T1	A		5	A							1							
B0Y5T1_AS	F		0	F							5							
PFC	U		8	U							8					1	1	1
Endosomal	B	B	5	A					2	2						6	4	6
cargo	—	0	.	—	0				5	9						8	4	1
receptor	0	Y	C	4	.	2			.	3						4	8	6
(Erp5),	6	5	A	G	9	0	2	8	3							4	2	2
putative	4	T	n	D	0	0	.	3	6	.						0	0	0
OS=Neosart	4	1	o	A	7	4	8	1	3	3	3	3	3	3	0	0	0	0

binding	B R	5 A 1 3	5 1 6	8 4 3
protein,	_ G	. _ 1	1 5 6	7 4 5
putative	0 2	C 1	4 9 6	0 0 0
OS=Neosart	0	A G	0 6	0 0
orya	2	D 0	6	
fumigata	3	A 1	6	
(strain	6	F 9	7	
CEA10 / CBS	0	U 8		
144.89 /		B 0		
FGSC		P		
A1163)		0		
GN=AFUB_0		0		
02360 PE=4		0		
SV=1		0		
		0		
		2		
		3		
		1		
		5		
		0		
		8		
		5		
tr B0XX33		.		
B0XX33_AS		C		
PFC FAD-		A		
dependent		D		
oxygenase,		A		
putative		F A		
OS=Neosart	A	U F		
orya	F	B U	9 5	
fumigata	U	P A	2 .	
(strain	B	0 _	1 6	
CEA10 / CBS	_	0 3	0 6	
144.89 /	0 B	0 G	5 2 6	8 9 9
FGSC	4 0	0 0 0	5 3 6	3 9 2
A1163)	7 X	4 0 . 1	. . 6	9 3 9
GN=AFUB_0	5 X	6 8 5 7	5 0 3 6	1 9 9
47560 PE=4	6 3 n	6 4 8 .	0 2 3 6	4 5 8
SV=1	0 3 o	2 0 8 6	7 6 3 7	5 5 7 0 0 0 0 0 0 0 0 0
tr B0Y002	A	5 A	8	
B0Y002_AS	F	0 F	6	
PFC Endo-	U B	8 U	3 9	7 9 8
1,3(4)-beta-	B 0	5 A 0	1 8	8 6 5
glucanase,	_ Y	. _ . 2	. 8	9 9 0
putative	0 0	C 3 8 1	2 0 6	6 9 0
OS=Neosart	4 0 n	A G 9 .	8 8 .	5 7 4
orya	5 2 o	D 0 6 4	5 7 6	3 3 3 3 0 0 0 0 0 0 0 0 0

fumigata	1		A	3				6										
(strain	4		F	0				7										
CEA10 / CBS	0		U	8														
144.89 /			B	0														
FGSC			P															
A1163)			0															
GN=AFUB_0			0															
45140 PE=4			0															
SV=1			0															
			4															
			4															
			2															
			8															
			5															
			0															
			8															
			5															
			0															
tr BOXZB5			.															
BOXZB5_AS			C															
PFC Ser/Thr			A															
protein			D															
phosphatas			A															
e family			F	A														
OS=Neosart	A		U	F														
orya	F		B	U					4									
fumigata	U		P	A					.									
(strain	B		0	_					3									
CEA10 / CBS	_		0	3					3									
144.89 /	0	B	0	G				7	8	3				7	8	9		
FGSC	4	0	0	0	0			1	6	3				6	6	5		
A1163)	3	X	4	4	.	1		.	0	3				1	1	7		
GN=AFUB_0	8	Z	y	2	1	5	4	6	2	2	3			9	0	6		
43810 PE=4	1	B	e	9	6	1	.	2	6	3	3			6	6	7		
SV=1	0	5	s	9	0	4	4	6	2	0	3	5	4	4	0	0	0	0
tr BOXQR8			5															
BOXQR8_AS			0	A														
PFC	A		8	F														
Endopolyph	F		5	U							2							
osphatase	U		.	A							.							
OS=Neosart	B		C	_							6							
orya	_		A	1							6				1			
fumigata	0	B	D	G				7	8	6				5	0	9		
(strain	1	0	A	1	0			6	4	6				2	5	6		
CEA10 / CBS	0	X	F	1	.			.	6	6				5	3	1		
144.89 /	9	Q	y	U	4	7	3	6	2	9	6			7	6	4		
FGSC	2	R	e	B	9	8	.	6	0	3	6			2	0	7		
A1163)	0	8	s	P	0	2	4	8	6	0	7	2	3	3	0	0	0	0

GN=AFUB_0		0																					
10920 PE=4		0																					
SV=1		0																					
		0																					
		1																					
		0																					
		7																					
		2																					
		5																					
		0																					
		8																					
		5																					
		.																					
tr BOXUQ5		C																					
BOXUQ5_AS		A																					
PFC		D																					
Allergen		A																					
Asp F4		F A																					
OS=Neosart	A	U F																					
orya	F	B U							7	2													
fumigata	U	P A							8	.													
(strain	B	0 _							4	6													
CEA10 / CBS	_	0 2							4	6										1			
144.89 /	0 B	0 G							3	4	6									3	2	7	
FGSC	2 0	0 0 0							4	3	6										2	6	6
A1163)	0 X	2 3 . 2							.	.	6										6	1	5
GN=AFUB_0	9 U y	0 8 5 5	3	0	3	6															0	4	9
20900 PE=4	0 Q e	3 3 0 .	2	9	3	6															1	0	2
SV=1	0 5 s	9 0 1 2 2	3	3	7	1	4	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
tr BOYCL1		5																					
BOYCL1_ASP		0																					
FC		8																					
Glyoxylase		5																					
family		.																					
protein,		C																					
putative		A A																					
OS=Neosart	A	D F																					
orya	F	A U							7	2													
fumigata	U	F A							4	.													
(strain	B	U _							6	6													
CEA10 / CBS	_	B 7							2	6													
144.89 /	0 B	P G							2	3	6										5	9	7
FGSC	9 0	0 0 0							5	6	6										4	2	7
A1163)	0 Y	0 5 . 1							.	.	6										6	0	1
GN=AFUB_0	5 C y	0 0 5 4	2	2	6	6															0	7	9
90580 PE=4	8 L e	0 1 6 .	2	7	6	6															7	4	0
SV=1	0 1 s	8 5 1 7 5	3	7	7	2	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

8
 0
 6
 5
 0
 8
 5
 .
 C
 tr|BOY2H6| A
 BOY2H6_AS D
 PFC Lipase, A
 putative F A
 OS=Neosart A U F
 orya F B U 7
 fumigata U P A 3
 (strain B 0 _ 1
 CEA10 / CBS _ 0 5 3
 144.89 / 0 B 0 G 5 6 8 5 7
 FGSC 6 0 0 1 0 1 3 8 6 4
 A1163) 1 Y 6 4 . . 2 1 9
 GN=AFUB_0 8 2 0 1 6 4 9 3 2 9 8
 61840 PE=4 4 H n 4 5 2 7 0 3 8 4 7
 SV=1 0 6 o 0 0 8 4 0 7 3 1 1 1 1 0 0 0 0 0 0 0 0 0 0

5
 0
 8
 5
 .
 C
 tr|BOYA75| A
 BOYA75_AS A
 PFC D
 Uncharacter A
 ized protein F A
 OS=Neosart A U F
 orya F B U 6 1
 fumigata U P A 8 .
 (strain B 0 _ 9 6
 CEA10 / CBS _ 0 8 5 6
 144.89 / 0 B 0 G 4 9 6 7 7 4
 FGSC 8 0 0 0 0 5 6 6 9 8 8
 A1163) 3 Y 8 3 . . 6 4 7 6
 GN=AFUB_0 6 A y 1 9 5 9 4 0 6 6 3 6 8
 83670 PE=4 7 7 e 4 0 6 . 1 9 6 6 7 0 2
 SV=1 0 5 s 4 5 5 6 5 7 7 7 2 2 1 0 0 0 0 0 0 0 0 0 0
 tr|BOXTX2| A B y 5 A 0 6 2 2 6 3 8 7
 BOXTX2_AS F 0 e 0 F . . 3 6 7 1 1 1 1 0 0 0 9 3 8 0 0 0

PFC Cortical	U X s	8	U 8 7 9	. 0																7 3 1
patch	B T	5	A 2			6 7														9 2 0
protein	_ X	.	_ 1			0 2														1 1 5
SUR7,	0 2	C 2				9 3														0 0 0
putative	1	A G				.														
OS=Neosart	9	D 0				3														
orya	4	A 2				3														
fumigata	0	F 3				3														
(strain	0	U 1																		
CEA10 / CBS		B 0																		
144.89 /		P																		
FGSC		0																		
A1163)		0																		
GN=AFUB_0		0																		
19400 PE=4		0																		
SV=1		1																		
		8																		
		9																		
		3																		
		5																		
		0																		
		8																		
tr B0Y2S3		5																		
B0Y2S3_AS																				
PFC		.																		
Endosomal		C																		
cargo		A																		
receptor		D																		
(Erp3),		A																		
putative		F A																		
OS=Neosart	A	U F																		
orya	F	B U					6 2													
fumigata	U	P A					4 .													
(strain	B	0 _					1 6													
CEA10 / CBS	_	0 5					9 6													
144.89 /	0 B	0 G					2 7 6													5 7 6
FGSC	5 0	0 0 0					7 6 6													5 6 0
A1163)	1 Y	5 3 . 3					. . 6													7 7 1
GN=AFUB_0	7 2	0 2 8 1 2 1 6 6																		1 6 1
51770 PE=3	7 S n	6 6 1 . 4 5 6 6																		5 3 5
SV=1	0 3 o	8 0 1 5 1 9 7 7					3 3 2 0 0 0 0 0 0 0 0 0													0 0 0
tr B0XQ77	A	5 A					6													
B0XQ77_AS	F B	0 F					1 3													6 5 6
PFC Dynein	U 0	8 U 0					5 9													5 9 6
light chain	B X	5 A . 2					. 1													9 7 0
(Tctex1),	_ Q y	. _ 5 1 1 3 8																		4 6 5
putative	0 7 e	C 1 2 . 4 2 6																		0 2 4
OS=Neosart	0 7 s	A G 6 7 3 5 . 2 2 2 2 0 0 0 0 0 0 0 0																		0 0 0

orya	9	D	0	6																
fumigata	0	A	9	6																
(strain	0	F	5	7																
CEA10 / CBS	0	U	5																	
144.89 /		B	0																	
FGSC		P																		
A1163)		0																		
GN=AFUB_0		0																		
09000 PE=4		0																		
SV=1		0																		
		0																		
		8																		
		8																		
		1																		
		5																		
		0																		
tr BOXQR6		8																		
BOXQR6_AS		5																		
PFC		.																		
Endosomal		C																		
cargo		A																		
receptor		D																		
(P24),		A																		
putative		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U																	
fumigata	U	P	A																	
(strain	B	0																		
CEA10 / CBS	—	0	1																	
144.89 /	0	B	0	G	2	5						8	3							
FGSC	1	0	0	1	0	5	9					7	0							
A1163)	0	X	1	1	.	2	.	0				2	9							
GN=AFUB_0	9	Q	y	0	4	7	7	2	0	8		5	2							
10900 PE=3	0	R	e	7	7	8	.	1	5	8		4	3							
SV=1	0	6	s	0	0	3	1	8	6	5	2	3	1	2	0	0	0	0	0	0
tr BOY4P0		5	A																	
BOY4P0_AS	A	0	F																	
PFC	F	8	U							4										
Calnexin	U	5	A							.										
OS=Neosart	B	.	—							6										
orya	—	C	4							6										
fumigata	0	B	A	G	6	5	6					5	6	5						
(strain	6	0	D	1	0	1	7	6				8	2	2						
CEA10 / CBS	9	Y	A	2	.	.	6	6				3	5	1						
144.89 /	7	4	F	8	5	5	8	7	6			3	6	4						
FGSC	6	P	n	U	5	3	2	6	5	9	6	3	0	4						
A1163)	0	0	o	B	0	6	5	3	3	0	7	5	5	4	0	0	0	0	0	0

																					3
																					7
																					4
																					4
																					5
																					0
																					8
																					5
tr BOY9D8																					.
BOY9D8_AS																					C
PFC Beta-																					A
1,6-																					D
glucanase,																					A
putative																					F A
OS=Neosart	A																				U F
orya	F																				B U
fumigata	U																				P A
(strain	B																				0 _
CEA10 / CBS	_																				0 8
144.89 /	0 B																				0 G
FGSC	8 0																				5 6
A1163)	0 Y																				6
GN=AFUB_0	6 9																				5 6
80680 PE=3	8 D n																				7 4 3
SV=1	0 8 o																				8 7 3
sp BOXRS8																					6
ECM14 ASP																					6
FC Putative																					5 6
metallocarb																					7 4 3
oxypeptidas																					8 7 3
e ecm14																					6 7 6
OS=Neosart																					2 1 2
orya																					0 9 1
fumigata																					0 8 0
(strain																					7 0 2 6 8 3 0 7 3 3 2 0 0 0 0 0 0 0 0 0
CEA10 / CBS	B																				5 2
144.89 /	e 0																				2 .
FGSC	c X																				8 3
A1163)	m R y																				5 3
GN=ecm14	1 S e																				6 0 3
PE=3 SV=1	4 8 s																				6 3 3
sp BOXZV8	B																				6 3 3
PLB3 ASPFC	0																				6 5 4
Lysophosph	p X																				3 2 2 0 0 0 0 0 0 0 0 0
olipase 3	l Z																				7 2 .
OS=Neosart	b V n																				3 4 0
orya	3 8 o																				3 3 1
																					5 3
																					6 4 8 3
																					5 0 0
																					0 4 2
																					3 5 4 4 0 0 0 0 0 0 0 0 0

fumigata																				. 3
(strain																				3 3
CEA10 / CBS																				3 3
144.89 /																				3 3
FGSC																				
A1163)																				
GN=plb3																				
PE=1 SV=1																				
																				5
																				0
																				8
																				5
																				.
tr BOXSZ4																				C
BOXSZ4_ASP																				A
FC																				D
Uncharacter																				A
ized protein																				F A
OS=Neosart	A																			U F
orya	F																			B U 5
fumigata	U																			P A 0
(strain	B																			0 _ 4
CEA10 / CBS	_																			0 2 5
144.89 /	0 B																			0 G 1 4 4 4 5
FGSC	2 0																			0 1 0 9 6 9 6 5
A1163)	8 X																			2 2 . 1 . . 1 6 5
GN=AFUB_0	3 S																			7 6 8 4 1 5 6 5 1 9
28320 PE=4	2 Z n																			7 8 4 . 8 8 6 7 7 0
SV=1	0 4 o																			1 0 2 1 5 5 7 2 2 2 2 0 0 0 0 0 0 0 0 0
																				5
tr BOXT37																				0
BOXT37_AS																				8
PFC Purine																				5
nucleoside																				.
permease,																				C
putative																				A A
OS=Neosart	A																			D F
orya	F																			A U 4 1
fumigata	U																			F A 6 .
(strain	B																			U _ 5 6
CEA10 / CBS	_																			B 2 4 6
144.89 /	0 B																			P G 4 9 6 4 7 1
FGSC	1 0																			0 0 0 3 6 6 7 4 7
A1163)	7 X																			0 0 . . . 6 3 9 3
GN=AFUB_0	8 T																			0 8 8 9 4 3 6 6 5 1 8
17890 PE=4	9 3 n																			0 1 3 . 0 2 6 6 0 6 3
SV=1	0 7 o																			1 0 1 9 3 9 7 7 2 2 1 0 0 0 0 0 0 0 0 0 0

```

7
4
5
5
0
8
5
tr|B0XNV1| .
B0XNV1_AS C
PFC GPI A
anchored D
protein, A
putative F A
OS=Neosart A U F
orya F B U 3
fumigata U P A 7
(strain B 0 _ 9
CEA10 / CBS _ 0 1 1
144.89 / 0 B 0 G 5 4 5 2 2
FGSC 1 0 0 1 0 8 6 9 6 7
A1163) 5 X 1 6 . . . 6 2 8
GN=AFUB_0 7 N 5 4 5 3 5 5 6 6 5 2
15760 PE=4 6 V n 4 2 7 . 2 2 6 3 5 6
SV=1 0 1 o 5 0 6 1 3 8 7 1 1 1 1 0 0 0 0 0 0 0 0 0 0 0

```

```

5
0
8
5
tr|B0YBD2| C
B0YBD2_AS A
PFC L-amino D
acid oxidase A
LaoA F A
OS=Neosart A U F
orya F B U
fumigata U P A
(strain B 0 _
CEA10 / CBS _ 0 7
144.89 / 0 B 0 G 7 3 3 4 3
FGSC 9 0 0 0 0 8 5 1 3 0
A1163) 2 Y 8 6 . 1 . 2 2 8 5
GN=AFUB_0 3 B 9 8 6 0 6 6 3 2 8 9
92390 PE=4 9 D n 8 1 5 . 9 1 4 3 1 8
SV=1 0 2 o 6 0 4 2 7 2 0 3 3 3 3 0 0 0 0 0 0 0 0 0 0 0
tr|B0Y449| A B y 5 A 0 1 4 5 3 2 3 2 3
B0Y449_AS F 0 e 0 F . 6 6 1 3 . 3 2 2 0 0 0 3 9 6 0 0 0

```


144.89 /

		5																		
		0																		
		8																		
		5																		
tr BOYEU1		.																		
BOYEU1_AS		C																		
PFC Toxin		A																		
biosynthesis		D																		
peroxidase,		A																		
putative		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U					0												
fumigata	U	P	A					.												
(strain	B	0	-					6												
CEA10 / CBS	-	0	4					6												
144.89 /	1	B	0	G				3	6										3	2
FGSC	0	0	0	0				3	1	6									9	3
A1163)	0	Y	9	2	0	1		6	1	6									2	0
GN=AFUB_1	3	E	7	7	.	1	3	.	5	6									2	7
00370 PE=4	7	U	n	5	8	8	.	2	9	2	6								9	5
SV=1	0	1	o	6	0	3	1	4	8	0	7	0	1	1	0	0	0	0	0	0

		5																		
		0																		
		8																		
		5																		
tr BOXN71		.																		
BOXN71_AS		C																		
PFC		A																		
Autophagy		D																		
protein		A																		
Atg27,		F	A																	
putative		U	F																	
OS=Neosart	A	B	U																	
orya	F	P	A																	
fumigata	U	0	-																	
(strain	B	0	1																	
CEA10 / CBS	-	0	1																	
144.89 /	0	B	0	G				3	3										3	2
FGSC	1	0	0	1	0			7	0										6	9
A1163)	4	X	1	5	.			.	7										0	8
GN=AFUB_0	8	N	y	4	2	7	8	3	3	0									6	2
14800 PE=4	0	7	e	5	5	5	.	3	7	3									3	1
SV=1	0	1	s	1	0	9	4	3	7	0	1	1	1	1	0	0	0	0	0	0
tr BOYF50	A	B	5	A	0	1		3	3	1									3	2
BOYF50_AS	F	0	0	F	.	1	2	0	0	.									4	6
PFC Acid	U	Y	n	8	U	7	.	9	.	4	6								7	1
phosphatas	B	F	o	5	A	4	9	3	4	1	6	1	2	2	0	0	0	6	4	2

e, putative	_	5	.	_	2		5	2	6							9	4	3
OS=Neosart	1	0	C	4			9	0	6							0	0	0
orya	0		A	G					6									
fumigata	1		D	0					6									
(strain	5		A	1					6									
CEA10 / CBS	7		F	0					7									
144.89 /	0		U	7														
FGSC			B	0														
A1163)			P															
GN=AFUB_1			0															
01570 PE=4			0															
SV=1			0															
			0															
			9															
			8															
			6															
			5															
			5															
			0															
			8															
			5															
tr B0XV35			.															
B0XV35_AS			C															
PFC			A															
Xylosidase/			D															
arabinoxidase,			A															
putative			F	A														
OS=Neosart	A		U	F														
orya	F		B	U					2									
fumigata	U		P	A					8									
(strain	B		0	_					9									
CEA10 / CBS	_		0	2					8									
144.89 /	0	B	0	G			4	9					2	3	2			
FGSC	2	0	0	0	0		0	3					8	1	7			
A1163)	1	X	2	4	.	1	.	.					5	0	4			
GN=AFUB_0	5	V	1	4	8	9	3	8	3				2	2	2			
21510 PE=3	1	3	n	0	8	0	.	7	0	3			0	2	6			
SV=1	0	5	o	0	0	6	1	1	3	3	2	2	2	2	0	0	0	0
tr B0XU21	A		5	A					0									
B0XU21_AS	F		0	F					.									
PFC	U		8	U					6									
Cutinase	B	B	5	A			3	2	6				3	1				
OS=Neosart	_	0	.	_	0		5	6	6				8	5				
orya	0	X	C	2	.		.	9	6				3	6				
fumigata	3	U	A	G	6	7	3	6	7	6			4	0				
(strain	0	2	n	D	1	6	.	6	7	4	6		7	1				
CEA10 / CBS	0	1	o	A	4	3	9	6	7	0	6	0	1	1	0	0	0	0

144.89 / 4 F 4 7
 FGSC 0 U 2
 A1163) B 0
 GN=AFUB_0 P
 30040 PE=3 0
 SV=1 0
 0
 0
 2
 9
 4
 3

sp|BOYED6|
 LAP1_ASPF
 C Leucine
 aminopepti
 dase 1
 OS=Neosart
 orya
 fumigata
 (strain
 CEA10 / CBS B 4 2 2 2 2
 144.89 / 0 0 3 6 7 7 5
 FGSC l Y # # . 1 . 7 3 5 2
 A1163) a E N N 8 3 3 1 1 4 4 4
 GN=lap1 p D n / / 3 . 8 2 1 2 5 6
 PE=3 SV=2 1 6 o A A 6 9 8 7 0 2 2 2 2 0 0 0 0 0 0 0 0 0 0 0
 5
 0
 8
 5

tr|BOYAM7
 |BOYAM7_A .
 SPFC Class V C
 chitinase A
 ChiB1 D
 OS=Neosart A F F
 orya F U U 0
 fumigata U B A .
 (strain B P _ 6
 CEA10 / CBS _ 0 8 6
 144.89 / 0 B 0 G 4 2 6 1 3
 FGSC 8 0 0 0 0 7 4 6 5 3
 A1163) 5 Y 0 1 . . 5 6 7 3
 GN=AFUB_0 2 A y 8 4 7 4 6 5 6 9 0
 85200 PE=3 0 M e 2 1 6 3 2 0 6 3 7
 SV=1 0 7 s 9 0 4 9 3 2 0 7 1 1 0 0 0 0 0 0 0 0 0 0 0 0

							6																						
							5																						
							0																						
tr BOXSP1							8																						
BOXSP1_AS							5																						
PFC Lectin							.																						
family							C																						
integral							A																						
membrane							D																						
protein,							A																						
putative							F	A																					
OS=Neosart	A						U	F																					
orya	F						B	U					2																
fumigata	U						P	A					4																
(strain	B						0	_					4																
CEA10 / CBS	_						0	2					8																
144.89 /	0	B					0	G				3	3						2	2	2								
FGSC	2	0					0	1	0				6	3						9	2	1							
A1163)	7	X					2	2	.	1			.	.						1	7	5							
GN=AFUB_0	8	S					7	1	7	6	3	1	3							0	5	8							
27870 PE=4	7	P	n				2	8	2	.	2	9	3							5	9	6							
SV=1	0	1	o				6	0	1	2	7	7	3	2	2	2	2	2	0	0	0	0	0	0	0	0	0	0	0
							5																						
							0																						
							8																						
							5																						
							.																						
tr BOY335							C																						
BOY335_AS							A																						
PFC							D																						
Uncharacter							A																						
ized protein							F	A																					
OS=Neosart	A						U	F																					
orya	F						B	U					2	2															
fumigata	U						P	A					2	.															
(strain	B						0	_					9	3															
CEA10 / CBS	_						0	5					4	3															
144.89 /	0	B					0	G				2	4	3										1	2	2			
FGSC	5	0					0	0	0				6	3	3									7	7	4			
A1163)	2	Y					5	4	.	1			.	.	3									1	4	2			
GN=AFUB_0	8	3	y				1	2	8	6	2	1	3	3										3	5	4			
52800 PE=4	0	3	e				6	8	3	.	4	1	3	3										2	9	2			
SV=1	0	5	s				9	0	2	7	5	5	3	3	2	3	2	0	0	0	0	0	0	0	0	0	0	0	0
tr BOXZX0	A	B					5	A	0			5	2											2	1	2			
BOXZX0_AS	F	0					0	F	.	7	4	1	1											8	3	2			
PFC	U	X	n				8	U	7	.	6	.	3											3	4	3			
Histidine	B	Z	o				5	A	4	1	2	2	7	2	2	2	2	2	0	0	0	0	5	4	4	0	0	0	0

acid	_ X	. _ 1	9 9																	0 0 9
phosphatas	0 0	C 3	5 6																	0 0 0
e, putative	3	A G	.																	
OS=Neosart	4	D 1	6																	
orya	6	A 4	6																	
fumigata	6	F 5	7																	
(strain	0	U 7																		
CEA10 / CBS		B 0																		
144.89 /		P																		
FGSC		0																		
A1163)		0																		
GN=AFUB_0		0																		
34660 PE=4		0																		
SV=1		3																		
		3																		
		9																		
		0																		
sp B0Y9Q9																				
BTGE_ASPF																				
C Probable																				
beta-																				
glucosidase																				
btgE																				
OS=Neosart																				2 1
orya																				0 .
fumigata																				0 3
(strain																				8 3
CEA10 / CBS	B																			5 0 3
144.89 /	0																			2 2 1
FGSC	b Y	# # 0																		8 3 3
A1163)	t 9	N N . 5 5 1 3 3																		2 9 0
GN=btgE	g Q n	/ / 4 . 6 4 3 3																		5 4 4
PE=3 SV=2	E 9 o	A A 9 8 5 7 3 3																		6 2 3
tr B0Y2Z9		5																		1 2 1 0 0 0 0 0 0 0 0 0
B0Y2Z9_AS		0																		
PFC		8 A																		
Mutanase	A	5 F																		
OS=Neosart	F	. U																		1
orya	U	C A																		.
fumigata	B	A _																		3
(strain	_	D 5																		3
CEA10 / CBS	0 B	A G																		4 1 3
144.89 /	5 0	F 0 0																		4
FGSC	2 Y	U 3 .																		9 9 3
A1163)	4 2	B 9 8																		3 3 5
GN=AFUB_0	4 Z n	P 4 9 1 4 8 1 3																		4 9 8 3
52440 PE=4	0 9 o	0 0 3 5 6 1 2 3																		8 1 3
																				5 9 9
																				2 1 1 0 0 0 0 0 5 1 0 0 0

SV=1		0																						
		0																						
		0																						
		5																						
		1																						
		3																						
		3																						
		5																						
		0																						
		8																						
		5																						
		.																						
tr BOY037		C																						
BOY037_AS		A																						
PFC		D																						
Uncharacter		A																						
ized protein		F																						
OS=Neosart	A	U																						
orya	F	B												1										
fumigata	U	P												8										
(strain	B	0												6										
CEA10 / CBS	_	0												0										
144.89 /	0 B	0											2	1							1	2	1	
FGSC	4 0	0											0	3	3							7	6	1
A1163)	5 Y	4	.									2	.	.								3	9	4
GN=AFUB_0	5 0	4	6									3	2	8	3							7	6	5
45510 PE=4	1 3 n	6	7	.								2	1	3								8	7	9
SV=1	0 7 o	3	0	9	6	9	7	3	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0

		5																					
		0																					
		8																					
		5																					
tr BOYCBO		.																					
BOYCBO_AS		C																					
PFC Lipase,		A																					
putative		D	A																				
OS=Neosart	A	A	F																				
orya	F	F	U											0									
fumigata	U	U	A											.									
(strain	B	B	_											6									
CEA10 / CBS	_	P	7											6									
144.89 /	0 B	0	G										4	1	6							1	2
FGSC	8 0	0	0	0									9	8	6							4	2
A1163)	9 Y	0	4	.									.	2	6							5	0
GN=AFUB_0	5 C	0	0	6								4	2	7	6							0	4
89550 PE=4	5 B n	8	2	4								6	5	5	6							4	6
SV=1	0 0 o	7	0	3	5	3	8	0	7	1	1	0	0	0	0	0	0	0	0	0	0	0	0

0
5
5
0
8
5
. C
tr|B0XML5| A
B0XML5_AS A
PFC L- D
asparaginas A
e F A
OS=Neosart A U F
orya F B U
fumigata U P A
(strain B 0 _
CEA10 / CBS _ 0 1
144.89 / 0 B 0 G 3 1 1 2 1
FGSC 0 0 0 0 0 9 8 2 5 6
A1163) 3 X 0 2 . . 1 2 5 6
GN=AFUB_0 1 M 3 7 7 4 3 7 2 1 0 4
03170 PE=3 7 L n 1 8 9 . 7 8 1 3 1 9
SV=1 0 5 o 0 0 3 7 9 4 0 1 1 1 1 0 0 0 0 0 0 0 0 0 0
sp|B0Y665|
PLB1_ASPFC
Lysophospholipase 1
OS=Neosart
orya
fumigata
(strain
CEA10 / CBS B 6 1 2 1 1
144.89 / 0 0 8 7 0 6 6
FGSC p Y # # . . 9 5 3 9
A1163) l 6 N N 7 4 6 1 2 2 2 1
GN=plb1 b 6 n / / 8 . 3 4 1 4 4 5
PE=1 SV=1 1 5 o A A 3 6 3 3 0 2 2 2 2 0 0 0 0 0 0 0 0 0 0
tr|B0XVA9| A 5 A 1
B0XVA9_AS F 0 F 6
PFC Lectin U 8 U 7
family B 5 A 9
integral _ B . _ 3 1 1 1
membrane 0 0 C 2 0 3 3 7 8 4
protein, 3 X A G . 1 6 . 6 2 4
putative 2 V y D 1 6 7 3 . 3 8 0 8
OS=Neosart 4 A e A 6 9 . 2 0 3 6 9 5
orya 7 9 s F 8 1 4 8 7 3 2 2 2 2 0 0 0 0 0 0 0 0 0 0

fumigata	0	U	0																
(strain		B	0																
CEA10 / CBS		P																	
144.89 /		0																	
FGSC		0																	
A1163)		0																	
GN=AFUB_0		0																	
32470 PE=4		3																	
SV=1		1																	
		8																	
		3																	
		5																	
		0																	
		8																	
		5																	
tr BOXWI1		.																	
BOXWI1_AS		C																	
PFC 1,3-		A																	
beta-		D																	
glucanosyltr		A																	
ansferase		F	A																
OS=Neosart	A	U	F																
orya	F	B	U																
fumigata	U	P	A																
(strain	B	0	-																
CEA10 / CBS	-	0	3																
144.89 /	0	B	0	G		4	1							1	1				
FGSC	3	0	0	1	0		9	3						0	6				
A1163)	6	X	3	3	.		.	5						5	4				
GN=AFUB_0	0	W	5	2	7	4	4	9	0					5	6				
36000 PE=3	0	I	n	2	0	1	.	6	3	9				4	5				
SV=1	0	1	o	2	0	5	1	0	3	5	1	1	1	1	0	0	0	0	0
tr BOXY72		5																	
BOXY72_AS		0																	
PFC Cell		8																	
wall		5	A																
glucanase,	A	.	F																
putative	F	C	U																
OS=Neosart	U	A	A																
orya	B	D	-																
fumigata	-	A	3																
(strain	0	B	F	G			1							1	1				
CEA10 / CBS	3	0	U	0	0		4	0						0	9	2			
144.89 /	9	X	B	9	.		0	7						0	8	3			
FGSC	8	Y	y	P	2	9	6	3	.	5				6	0	8			
A1163)	7	7	e	0	5	3	.	6	5	3				8	3	8			
GN=AFUB_0	0	2	s	0	0	3	3	3	9	3	1	1	1	1	0	0	0	0	0

39870 PE=4		0																						
SV=1		0																						
		3																						
		9																						
		0																						
		6																						
		5																						
		0																						
		8																						
		5																						
tr BOYEU4		.																						
BOYEU4_AS		C																						
PFC		A																						
Glycosyl		D																						
hydrolase,		A																						
putative		F	A																					
OS=Neosart	A	U	F																					
orya	F	B	U					3																
fumigata	U	P	A					.																
(strain	B	0	-					3																
CEA10 / CBS	-	0	4					3																
144.89 /	1	B	0	G			9	3														1		
FGSC	0	0	0	0	0		8	9	3					7	7	2								
A1163)	0	Y	9	0	.		.	3	3					3	9	6								
GN=AFUB_1	9	E	7	3	6	7	8	9	1	3				4	0	7								
00920 PE=4	2	Y	n	9	9	6	.	8	2	0	3			6	7	6								
SV=1	0	4	o	9	0	2	8	3	9	1	3	3	3	4	0	0	0	0	5	8	0	0	0	0

tr BOXUD4		5																					
BOXUD4_AS		.																					
PFC		C																					
Glycosyl		A																					
hydrolase,		D																					
putative		A	A																				
OS=Neosart	A	F	F																				
orya	F	U	U					8															
fumigata	U	B	A					4															
(strain	B	P	-					8															
CEA10 / CBS	-	0	2					6															
144.89 /	0	B	0	G			6	8															
FGSC	2	0	0	0	0		9	.						9	8	7							
A1163)	0	X	0	3	.		.	6						5	7	1							
GN=AFUB_0	3	U	1	2	8	2	6	3	6					7	6	2							
20340 PE=4	4	D	n	9	7	0	.	2	0	6				3	5	0							
SV=1	0	4	o	8	0	1	1	0	5	7	1	1	1	1	0	0	0	9	8	9	0	0	0

		7																
		5																
		0																
		8																
tr B0Y2U6		5																
B0Y2U6_AS		.																
PFC Alpha		C																
glucosidase		A																
II, alpha		D																
subunit,		A																
putative		F A																
OS=Neosart	A	U F																
orya	F	B U																
fumigata	U	P A																
(strain	B	0 _																
CEA10 / CBS	_	0 5																
144.89 /	0 B	0 G				1									1			
FGSC	5 0	0 0 0				0 8									1 2 9			
A1163)	2 Y	5 3 . 1				9 0									7 6 7			
GN=AFUB_0	0 2	0 5 7 0 9 . 1													2 3 0			
52020 PE=3	2 U n	9 0 7 . 6 6 8													2 1 3			
SV=1	0 6 o	1 0 1 5 7 5 9 2 3 1 2 0 0 0 0 0													0 3 4 0 0 0 0			
		5																
		0																
		8																
tr B0XU27		5																
B0XU27_AS		.																
PFC		C																
Oxidoreduct		A																
ase, FAD-		D																
binding,		A																
putative		F A																
OS=Neosart	A	U F																
orya	F	B U													0			
fumigata	U	P A													.			
(strain	B	0 _													6			
CEA10 / CBS	_	0 2													6			
144.89 /	0 B	0 G				5									6			
FGSC	3 0	0 1 0				0 6 6									7 6			
A1163)	0 X	2 4 .				. 8 6									4 1			
GN=AFUB_0	1 U	9 4 8 4 4 4 1 6													6 7			
30100 PE=4	0 2 n	4 8 8 . 7 9 9 6													9 0			
SV=1	0 7 o	9 0 5 2 3 4 7 7 0 1 1 0 0 0 0 0													0 2 2 0 0 0 0			
tr B0Y6J5 B	A B	5 A 0				7 3									4 3 3			
OY6J5_ASPF	F 0 y	0 F . 1 6 3 9													8 1 7			
C alpha-1,2-	U Y e	8 U 6 . 4 . 3													9 6 6			
Mannosidas	B 6 s	5 A 4 6 1 6 8 1 1 1 1 0 0 0 0 0													3 1 0 0 0 0 0			

```

e                _ J    . _ 4    0 4                9 0 4
OS=Neosart     0 5    C 4    6 .
orya           6    A G    3
fumigata       7    D 1    3
(strain        1    A 0    3
CEA10 / CBS    8    F 0    3
144.89 /      0    U 7
FGSC           B 0
A1163)        P
GN=AFUB_0     0
67180 PE=3    0
SV=1          0
              0
              6
              5
              3
              7
              5
              0
              8
tr|BOYOF9|     8
BOYOF9_AS     5
PFC           .
Alpha,alpha    C
-trehalose    A
glucohydrol   D
ase           A
TreA/Ath1     F A
OS=Neosart   A    U F
orya         F    B U    0
fumigata     U    P A    .
(strain      B    0 _    6
CEA10 / CBS  _    0 3    6
144.89 /    0 B    0 G    6
FGSC        4 0    0 0 0    2 6    3 2
A1163)     6 Y    4 2 .    1 9 6    0 9
GN=AFUB_0  0 0    5 2 8 1 0 1 9 6    3 4
46050 PE=4  5 F n  1 8 0 . 7 1 2 6    8 7
SV=1       0 9 o  7 0 7 2 2 7 6 7 0 1 1 0 0 0 0 0 2 0 0 0

```

Proteins with signal peptide which are quantified in 115146 and not in Ku80 at 10d and 2h of extraction (Ranked by average iBAQ intensity in 115146)	Statistics (rank from the high	Number of identified peptides specific to each protein in each sample	iBAQ (intensity-based absolute quantification) values

										est mea n of iBA Q valu es to the low est to rank fro m the mos t abu nda nt prot ein to the leas t)													
	P	U									M	U	U	U	U	U	U	i	i	i	i	i	i
	re	r	n								M	e	n	n	n	n	n	B	B	B	B	B	B
	s	o	i								e	a	i	i	i	i	i	A	A	A	A	A	A
	U	t	q								M	a	q	q	q	q	q	Q	Q	Q	Q	Q	Q
	n	u									n	U	u	u	u	u	u
	i	w	e	S	l						i	e	e	e	e	e	e	1	1	1	1	1	1
	p	i	.	e	.						i	B	q	p	p	p	p	0	0	0	0	0	0
	r	n	t	s	q						q	A	e	e	e	e	e	-	-	-	-	-	-
	o	g	h	e	u	w					Q	e	p	p	p	p	p	2	2	2	2	2	2
	t	_	B	q	e	e					P	t	i	i	i	i	i	h	h	h	h	h	h
	.	4	p	L	s	u	n	i			1	e	d	d	d	d	d	-	-	-	-	-	-
	a	6	r	A	i	e	c	g			0	p	e	e	e	e	e	1	1	1	D	D	D
	G	c	n	t	T	n	c	.	t		d	t	s	s	s	s	s	1	1	1	e	e	e
	e	e	d	e	_	a	e	l	.		1	d	1	1	1	1	1	5	5	5	l	l	l
	n	s	a	i	A	l	.	e	.		1	e	0	0	0	0	0	.	.	.	t	t	t
	e	s	b	n	f	c	n	k			5	s	d	d	d	d	d	1	1	1	a	a	a
	.	i	s	_	2	p	o	g	D		1	1	-	-	-	-	-	4	4	4	K	K	K
Fasta.headers	l	o	e	i	9	e	v	t	a		4	0	2	2	2	2	2	6	6	6	U	U	U
	D	n	nt	d	3	p	e	h	.		6	d	h	h	h	h	h	-	-	-	8	8	8

	in		t r		1							1	2	3	0	0	0
	K		i a		1												
	u		d g		5												
	8		e e		1												
	0		..		4												
	at		(..		6												
	3		S														
	0		c														
	d		o														
	2		r														
	h		e														
			o														
			f														
			S														
			i														
			g														
			n														
			a														
			l														
			P														
			i														
			f														
			p														
			r														
			e														
			d														
			i														
			c														
			t														
			e														
			d														
)														
tr BOYOH2	A		5	A													
BOYOH2_AS	F		0	F		7	8										
PFC	U		8	U		4	.										
Uncharacter	B		5	A		6	6										
ized protein	_		.	_		2	6					8	8	4			
OS=Neosart	0	B	C	3		2	6					6	7	9			
orya	4	0	A	G	0	3	6	6				4	8	5			
fumigata	6	Y	D	0	.	3	3	6	6			7	9	0			
(strain	1	0	A	2	8	0	2	.	.	6		8	0	0			
CEA10 / CBS	8	H	n	F	1	6	.	9	6	6	6	0	0	0			
144.89 /	0	2	o	U	9	9	3	7	1	7	7	9	9	8	0	0	0

FGSC		B	0															
A1163)		P																
GN=AFUB_0			0															
46180 PE=4			0															
SV=1			0															
			0															
			4															
			5															
			3															
			0															
			5															
			0															
			8															
tr BOY1D4			5															
BOY1D4_AS			.															
PFC			C															
Disulfide			A															
isomerase			D															
(TigA),			A															
putative			F	A														
OS=Neosart	A		U	F														
orya	F		B	U														
fumigata	U		P	A														
(strain	B		0	-														
CEA10 / CBS	-		0	5			4						4	5	4			
144.89 /	0	B	0	G			4	8					6	0	7			
FGSC	5	0	0	1	0			0	3				6	8	6			
A1163)	9	Y	5	2	.	5		.	7				2	5	5			
GN=AFUB_0	8	1	y	8	2	8	4	3	2	9			8	3	6			
59890 PE=3	9	D	e	5	6	6	.	6	5	0	1	1	1	1				0
SV=1	0	4	s	7	0	7	6	8	3	0	4	4	4	4	0	0	0	0
tr BOXUX8			5															
BOXUX8_AS			0															
PFC			8															
Translocon-			5															
associated			.	A														
protein,	A		C	F														
alpha	F		A	U														
subunit,	U		D	A														
putative	B		A	-														
OS=Neosart	-		F	2									4	1	7			
orya	0	B	U	G						2	3		2	8	0			
fumigata	3	0	B	1	0					8	9		4	5	9			
(strain	1	X	P	6	.	4				.	8		7	7	0			
CEA10 / CBS	8	U	y	0	1	8	2	2	1	4			3	1	8			
144.89 /	0	X	e	0	2	4	.	6	5	0			0	0	0			
FGSC	0	8	s	0	0	1	1	1	8	0	4	4	2	6	0	0	0	0

A1163)		P																		
GN=AFUB_0		0																		
38220 PE=4		0																		
SV=1		0																		
		0																		
		3																		
		7																		
		4																		
		4																		
		5																		
		0																		
		8																		
tr B0YBS2		5																		
B0YBS2_AS		.																		
PFC		C																		
Cysteine-		A																		
rich		D																		
secreted		A																		
protein		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U																	
fumigata	U	P	A																	
(strain	B	0	_																	
CEA10 / CBS	_	0	7																	
144.89 /	0	B	0	G																
FGSC	8	0	0	0	0															
A1163)	7	Y	8	1	.	2														
GN=AFUB_0	6	B	5	0	8	2	3	.												
87640 PE=4	4	S	n	1	6	1	.	4	8	0	6									
SV=1	0	2	o	7	0	1	4	3	6	0	7									
sp B0Y766																				
P20D1_ASP																				
FC Probable																				
carboxypept																				
idase																				
AFUB_0727																				
30	A																			
OS=Neosart	F																			
orya	U																			
fumigata	B																			
(strain	_																			
CEA10 / CBS	0	B																		
144.89 /	7	0																		
FGSC	2	Y																		
A1163)	7	7	y																	
GN=AFUB_0	3	6	e																	
72730 PE=3	0	6	s																	

BOY1U6_AS	F 0 e	0 F .	8 9 2 6 .											4 5 8
PFC	U Y s	8 U 6 .	1 . 3 3											5 7 6
Agmatinase, putative	B 1 _ U	5 A 7 4 . _ 8	1 1 3 7 6 3											9 4 0 2 9 9
OS=Neosart orya	0 6	C 5 A G	9 6 3 6 3											0 0 0 0 0 0
fumigata (strain)	0 8	D 1 A 3	. 6 3											
CEA10 / CBS 144.89 / FGSC A1163)	9 0	F 1 U 8 B 0	7 3											
GN=AFUB_0 60890 PE=3 SV=1														
tr BOXR6 BOXR6_AS														
PFC														
Endosomal cargo receptor (P24), putative														
OS=Neosart orya	A F	U B	F U											
fumigata (strain)	B U	P A												1 4 5 . 9 6
CEA10 / CBS 144.89 / FGSC A1163)	_ 0	0 1 B	5 6 2 1 6											1 1 1 4 7 4
GN=AFUB_0 10900 PE=3 SV=1	9 0	Q R e	y 7 7 8 . 1 5 3 6											6 6 2 0 0 0
tr BOYDZ9 BOYDZ9_AS	0 6	s 0 0	3 1 8 6 3 7											0 0 0 0 0 0 0
PFC Ser/Thr protein phosphatase family	A B F U B D _ Z 0	5 A 8 U 5 A 7 3 . _ 3 C	7 1 5 5 . 6 4 8 1 1 1 0 9 0 0 0											1 1 1 4 5 6 6 8 2 8 5 9 0 4 1 0 0 0 0 0 0

protein	9		A	G			3													0	0	0	
OS=Neosart	7		D	0			.																
orya	3		A	0			3																
fumigata	4		F	1			3																
(strain	0		U	6																			
CEA10 / CBS			B	0																			
144.89 /			P																				
FGSC			0																				
A1163)			0																				
GN=AFUB_0			0																				
97340 PE=4			0																				
SV=1			9																				
			4																				
			6																				
			4																				
			5																				
			0																				
			8																				
			5																				
tr BOYCQ5			.																				
BOYCQ5_AS			C																				
PFC SUN			A																				
domain			A																				
protein			D																				
(Uth1),			A																				
putative			F	A																			
OS=Neosart	A		U	F																			
orya	F		B	U					1	5													
fumigata	U		P	A					4	.													
(strain	B		0	-					7	6													
CEA10 / CBS	-		0	7					4	6										1	1		
144.89 /	0	B	0	G					4	4	6									6	9	7	
FGSC	9	0	0	0	0				3	6	6									6	7	8	
A1163)	1	Y	8	5	.	2			.	6	6									2	9	1	
GN=AFUB_0	0	C	y	8	4	4	3	4	5	.	6									1	8	5	
91030 PE=4	3	Q	e	5	5	9	.	1	0	6	6									0	0	0	
SV=1	0	5	s	0	0	6	2	4	4	7	7	5	5	7	0	0	0	0	0	0	0	0	0
tr BOYEP2	A		5	A																			
BOYEP2_AS	F		0	F					1	3													
PFC Cell	U		8	U					1	.													
wall serine-	B		5	A					1	6													
threonine-	-		.	-					9	6										1	1		
rich	0	B	C	4					2	0	6									3	9	0	
galactoman	9	0	A	G	0				7	9	6									2	4	8	
noprotein	9	Y	D	0	.	3			.	6	6									4	9	3	
Mp1	8	E	A	3	8	3	2	3	.	6										5	0	7	
OS=Neosart	8	P	n	F	2	2	.	8	6	6	6									0	9	0	
orya	0	2	o	U	4	5	5	4	1	7	7	4	3	4	0	0	0	0	0	0	0	0	0

subunit 1	2		F	7					3	3									
OS=Neosart	0		U	9															
orya			B	0															
fumigata			P																
(strain			0																
CEA10 / CBS			0																
144.89 /			0																
FGSC			0																
A1163)			2																
GN=AFUB_0			2																
22820 PE=3			2																
SV=1			9																
			5																
			0																
			8																
tr BOYCL1			5																
BOYCL1_ASP			.																
FC			C																
Glyoxylase			A																
family			D																
protein,			A																
putative			F	A															
OS=Neosart	A		U	F															
orya	F		B	U					6	1									
fumigata	U		P	A					7	.									
(strain	B		0	_					6	6									
CEA10 / CBS	_		0	7					8	6									
144.89 /	0	B	0	G					2	0	6			8	7	3			
FGSC	9	0	0	0					5	3	6			8	4	9			
A1163)	0	Y	8	5	.	1			.	.	6			6	4	9			
GN=AFUB_0	5	C	y	8	0	5	4	2	2	3	6			0	4	9			
90580 PE=4	8	L	e	0	1	6	.	2	7	3	6			2	2	7			
SV=1	0	1	s	6	5	1	7	5	3	3	7	2	2	1	0	0	0	0	0
sp BOXRV0			5																
DPP5_ASPF			0																
C			8	A															
Dipeptidyl-	A		5	F															
peptidase 5	F		.	U					5	7									
OS=Neosart	U		C	A					8	.									
orya	B		A	_					9	3									
fumigata	_		D	2					3	3									
(strain	0	B	A	G					7	9	3			6	4	7			
CEA10 / CBS	2	0	F	0	0				9	3	3			2	0	3			
144.89 /	4	X	U	9	.				.	.	3			7	6	4			
FGSC	9	R	B	0	7				7	7	3	3		6	2	3			
A1163)	2	V	n	P	3	3	1	2	4	3	3			8	0	0			
GN=AFUB_0	0	0	o	0	0	9	6	1	2	3	3	8	6	8	0	0	0	0	0

3
1
5
0
8
5
tr|B0XTX2|
B0XTX2_AS
PFC Cortical
patch
protein
SUR7,
putative
OS=Neosart A U F A
orya F B U
fumigata U P A
(strain B 0 _
CEA10 / CBS _ 0 2
144.89 / 0 B 0 G 2 5 4 3 7
FGSC 1 0 0 0 0 6 1 8 3 2
A1163) 9 X 1 2 . . 6 6 9 3
GN=AFUB_0 4 T y 8 3 8 6 2 6 6 7 2 9
19400 PE=4 0 X e 9 1 2 . 3 0 3 1 4 4
SV=1 0 2 s 3 0 1 7 9 9 0 1 1 1 1 0 0 0 0 0 0 0 0 0 0
5
0
8
5
tr|B0XY72|
B0XY72_AS
PFC Cell
wall
glucanase,
putative
OS=Neosart A U F A
orya F B U
fumigata U P A
(strain B 0 _
CEA10 / CBS _ 0 3
144.89 / 0 B 0 G 5 4 6 4
FGSC 3 0 0 0 0 4 1 6 0 6
A1163) 9 X 3 9 . 0 4 8 7 7
GN=AFUB_0 8 Y y 9 2 9 6 3 . 6 5 5 9
39870 PE=4 7 7 e 0 5 3 . 6 5 5 0 0 5
SV=1 0 2 s 6 0 3 3 3 9 0 2 2 2 2 0 0 0 0 0 0 0 0 0
tr|B0Y5L1|
B0Y5L1_ASP F 0 n 0 F . . 2 2 8 . 4 6 3
FC GPI U Y o 8 U 8 9 1 . 5 6 2 2 1 0 0 0 2 8 5 0 0 0

anchored	B 5	5 A 0	6 6 6	5 6 6
cell wall	_ L	. _ 7	2 2 6	9 7 0
protein,	0 1	C 4	0 6	0 0 0
putative	6	A G	6	
OS=Neosart	3	D 0	6	
orya	7	A 6	6	
fumigata	7	F 7	7	
(strain	0	U 0		
CEA10 / CBS		B 0		
144.89 /		P		
FGSC		0		
A1163)		0		
GN=AFUB_0		0		
63770 PE=4		0		
SV=1		6		
		2		
		0		
		3		
		5		
		0		
		8		
		5		
tr B0XZB5		.		
B0XZB5_AS		C		
PFC Ser/Thr		A		
protein		D		
phosphatas		A		
e family		F A		
OS=Neosart	A	U F		
orya	F	B U	1	
fumigata	U	P A	.	
(strain	B	0 _	3	
CEA10 / CBS	_	0 3	3	
144.89 /	0 B	0 G	7 3 3	5 2 3
FGSC	4 0	0 0 0	1 7 3	5 4 3
A1163)	3 X	4 4 . 1	. 7 3	6 5 1
GN=AFUB_0	8 Z y	2 1 5 4	6 2 7 3	6 3 0
43810 PE=4	1 B e	9 6 1 .	2 6 0 3	9 6 5
SV=1	0 5 s	9 0 4 4	6 2 0 3	2 1 1 0 0 0 0 0 0 0 0 0
tr B0XTT1	A	5 A	3	
B0XTT1_AS	F	0 F	7	
PFC	U B	8 U	6 1	3 2 4
Mitochondri	B 0	5 A 0	8 0	9 6 5
al dihydroxy	_ X	. _ . 1	. 2	3 8 1
acid	0 T	C 2 5 2	6 2 6	2 6 1
dehydratase	2 T n	A G 0 .	4 5 .	9 4 5
, putative	9 1 o	D 1 7 6	2 6 6	5 5 5 5 0 0 0 0 0 0 0 0 0


```

FGSC 0
A1163) 0
GN=AFUB_0 0
25060 PE=4 0
SV=1 2
4
4
9

sp|BOXRS8|
ECM14_ASP
FC Putative
metallocarb
oxyptidas
e ecm14
OS=Neosart 0
orya .
fumigata 6
(strain 6
CEA10 / CBS B 6 3 6 2 3
144.89 / e 0 0 6 0 6 6 5
FGSC c X # # . . 8 6 2 4
A1163) m R y N N 9 6 5 0 8 6 8 7
GN=ecm14 1 S e / / 5 . 8 6 1 6 4 8
PE=3 SV=1 4 8 s A A 1 7 6 8 0 7 1 1 0 0 0 0 0 0 0 0 0 0 0
5
0
8
5
tr|BOXN52|
BOXN52_AS .
PFC Hsp70 C
family A
chaperone D
Lhs1/Orp15 A
0, putative F A
OS=Neosart A U F
orya F B U 5
fumigata U P A .
(strain B 0 _ 3
CEA10 / CBS _ 0 1 3
144.89 / 0 B 0 G 1 3 3 3 3 2
FGSC 1 0 0 1 0 0 0 3 2 2 5
A1163) 4 X 1 5 . 8 0 3 3 4 2
GN=AFUB_0 6 N 4 0 5 9 . 0 3 1 7 1
14610 PE=3 1 5 n 3 5 8 1 9 7 4 3 8 9 5
SV=1 0 2 o 2 0 3 0 7 5 0 3 6 6 4 0 0 0 0 0 0 0 0 0 0
tr|BOXQR8| A B y 5 A 0 3 6 7 2 2 2 4
BOXQR8_AS F 0 e 0 F . . 6 6 9 2 2 2 2 0 0 0 5 0 2 0 0 0

```

PFC	U X s	8	U 7 4 8	.	5							6	8	0
Endopolyph	B Q	5	A 8		2 4							9	6	7
osphatase	_ R	.	_ 2		0 7							9	9	4
OS=Neosart	0 8	C	1		6 3							0	0	0
orya	1	A	G		.									
fumigata	0	D	1		3									
(strain	9	A	1		3									
CEA10 / CBS	2	F	4		3									
144.89 /	0	U	9											
FGSC		B	0											
A1163)		P												
GN=AFUB_0		0												
10920 PE=4		0												
SV=1		0												
		0												
		1												
		0												
		7												
		2												
		5												
		0												
		8												
		5												
		.												
tr BOY335		C												
BOY335_AS		A												
PFC		D												
Uncharacter		A												
ized protein		F	A											
OS=Neosart	A	U	F											
orya	F	B	U		2 2									
fumigata	U	P	A		9 .									
(strain	B	0	_		3 3									
CEA10 / CBS	_	0	5		0 3									
144.89 /	0 B	0	G		2 2 3						2	2	3	
FGSC	5 0	0	0 0		6 6 3						6	4	6	
A1163)	2 Y	5	4 . 1		. . 3						3	6	9	
GN=AFUB_0	8 3 y	1	2 8 6 2 1 6 3								5	5	0	
52800 PE=4	0 3 e	6	8 3 . 4 1 6 3								0	0	8	
SV=1	0 5 s	9	0 2 7 5 5 7 3			2	2	3	0	0	0	0	0	0
tr BOXW28	A	5	A		0									
BOXW28_AS	F B	0	F		3 2 .						2	2		
PFC	U 0	8	U 0		5 6 6						8	3		
Oligosaccha	B X	5	A .		. 0 6						2	8		
ryl	_ W	.	_ 6 7 3 9 6 6								8	4		
transferase	0 2 n	C	2 9 . 2 3 4 6								4	4		
subunit	2 8 o	A	G 1 1 5 9 0 6			1	1	0	0	0	0	0	0	0

(Gamma),	3		D	0																	6	
putative	3		A	6																		6
OS=Neosart	6		F	2																		7
orya	0		U	8																		
fumigata			B	0																		
(strain			P																			
CEA10 / CBS				0																		
144.89 /				0																		
FGSC				0																		
A1163)				0																		
GN=AFUB_0				2																		
23360 PE=4				2																		
SV=1				7																		
				9																		
				5																		
				0																		
				8																		
				5																		
			.																			
tr B0Y479			C																			
B0Y479_AS			A																			
PFC			D																			
Uncharacter			A																			
ized protein			F	A																		
OS=Neosart	A		U	F																		
orya	F		B	U																		2
fumigata	U		P	A																		2
(strain	B		0																			3
CEA10 / CBS	_		0	5																		9
144.89 /	0	B	0	G						1									2	2	2	
FGSC	5	0	0	0	0					0									3	2	2	
A1163)	6	Y	5	9	.					3	.								9	2	0	
GN=AFUB_0	8	4	5	2	8	6	9	.		3									3	4	0	
56810 PE=4	1	7	n	6	7	7	.	5	5	3									0	6	1	
SV=1	0	9	o	3	0	2	2	2	5	3	4	4	4	4	0	0	0	0	0	0	0	0
tr B0Y4T7			5	A																		
B0Y4T7_ASP	A		0	F																		
FC DUF907	F		8	U																		1
domain	U		5	A																		7
protein	B	.	.	_																		8
OS=Neosart	_		C	4																		2
orya	0	B	A	G						7	7									1	2	1
fumigata	7	0	D	1	0					8	6									6	1	5
(strain	0	Y	A	3	.					.	.									4	2	8
CEA10 / CBS	2	4	F	3	9	3	7	5	6											5	2	0
144.89 /	6	T	n	U	4	2	.	2	3	6										6	1	6
FGSC	0	7	o	B	0	8	9	1	8	7	2	2	2	2	0	0	0	0	0	0	0	0

A1163)		P																		
GN=AFUB_0		0																		
70260 PE=4		0																		
SV=1		0																		
		0																		
		6																		
		8																		
		4																		
		3																		
tr BOY269																				
BOY269_AS																				
PFC																				
NlpC/P60-																				
like cell-wall																				
peptidase,																				
putative																				
OS=Neosart		5																		
orya		0																		
fumigata		8																		
(strain		5																		
CEA10 / CBS		.																		
144.89 /		C																		
FGSC		A																		
A1163)		D																		
GN=AFUB_0		A																		
61470 PE=4		F	A																	
SV=1;tr BOY	A	U	F																	
9E0 BOY9E0	F	B	U					1	1											
_ASPFC	U	P	A					6	.											
Uncharacter	B	0	_					9	3											
ized protein	_	0	5					8	3											
OS=Neosart	0	B	0	G				3	9	3					1	1	1			
orya	6	0	0	1	0			8	6	3					4	8	7			
fumigata	1	Y	6	3	.	1		.	.	3					8	6	5			
(strain	4	2	y	0	7	8	2	3	7	6	3				2	0	3			
CEA10 / CBS	7	6	e	0	3	0	.	5	6	6	3				9	4	6			
144.89 /	0	9	s	4	0	9	3	9	3	7	3	1	1	2	0	0	0	0	0	0
tr BOYAM7	A		5	A						1										
BOYAM7_AS	F		0	F						5										
PFC Class V	U		8	U						9										
chitinase	B		5	A						9										
ChiB1	_	B	.	_				4	8						1	2	1			
OS=Neosart	0	0	C	8	0			7	6						4	0	3			
orya	8	Y	A	G	.			.	.						0	3	6			
fumigata	5	A	y	D	0	7		4	6	6					2	0	6			
(strain	2	M	e	A	1	6		3	2	6					5	4	7			
CEA10 / CBS	0	7	s	F	4	4	9	3	2	7	1	1	1	1	0	0	0	0	0	0

144.89 / 0 U 1
FGSC B 0
A1163) P
GN=AFUB_0 0
85200 PE=3 0
SV=1 0
0
8
2
9
6
5
0
8
5
tr|BOY7N9| .
BOY7N9_AS C
PFC Cell A
wall D
glucanase, A
putative F A
OS=Neosart A U F
orya F B U 1
fumigata U P A 5
(strain B 0 _ 2
CEA10 / CBS _ 0 6 8
144.89 / 0 B 0 G 4 0 1 1 1
FGSC 7 0 0 0 0 5 3 4 3 7
A1163) 4 Y 7 8 . 1 . . 6 6 5
GN=AFUB_0 4 7 2 5 7 0 4 6 3 4 0 9
74470 PE=4 7 N n 5 1 8 . 3 7 3 0 3 8
SV=1 0 9 o 4 0 7 3 6 4 3 1 1 1 1 0 0 0 0 0 0 0 0 0 0
tr|BOXP24| 5
BOXP24_AS 0
PFC 8
Disulfide 5 A
isomerase, A . F
putative F C U 1 2
OS=Neosart U A A 4 .
orya B D _ 6 3 2
fumigata _ A 1 3 3 .
(strain 0 B F G 5 2 3 0 1
CEA10 / CBS 0 0 U 0 0 1 1 3 0 4 9
144.89 / 5 X B 5 . . . 3 E 0 8
FGSC 6 P P 3 8 7 4 2 6 3 + 5 4
A1163) 7 2 n 0 2 6 . 7 9 6 3 0 4 2
GN=AFUB_0 0 4 o 0 0 8 5 8 9 7 3 3 2 2 0 0 0 0 5 0 5 0 0 0

05670 PE=4	0																		
SV=1	0																		
	0																		
	5																		
	5																		
	7																		
	5																		
	0																		
	8																		
	5																		
	.																		
tr B0XUQ5	C																		
B0XUQ5_AS	A																		
PFC	D																		
Allergen	A																		
Asp F4	F	A																	
OS=Neosart	A	U	F																
orya	F	B	U		1	1													
fumigata	U	P	A		3	.													
(strain	B	0	—		8	3													
CEA10 / CBS	—	0	2		8	3													
144.89 /	0	B	0	G	3	3	3					2							
FGSC	2	0	0	0	0	4	6	3				1	9	9					
A1163)	0	X	2	3	.	2	.	.	3			9	8	8					
GN=AFUB_0	9	U	y	0	8	5	5	3	0	6	3				7	0	7		
20900 PE=4	0	Q	e	3	3	0	.	2	9	6	3				6	5	0		
SV=1	0	5	s	9	0	1	2	2	3	7	3	2	1	1	0	0	0	0	0
sp B0Y1V8																			
PEPA_ASPF																			
C Aspartic																			
protease																			
pep1																			
OS=Neosart												0							
orya												.							
fumigata												6							
(strain												6							
CEA10 / CBS	B					4	1	6							1	1			
144.89 /	0					0			1	3	6				1	5			
FGSC	p	Y	#	#	.	3	.	3	6			5	0						
A1163)	e	1	N	N	6	3	3	6	2	6					6	8			
GN=pep1	p	V	n	/	/	8	.	9	1	4	6				2	7			
PE=3 SV=1	1	8	o	A	A	8	2	5	3	5	7	1	1	0	0	0	0	0	0
tr B0XN71	A	B		5	A	0			3	1	1				6	2	8		
B0XN71_AS	F	0		0	F	.			7	2	.				9	0	8		
PFC	U	X	y	8	U	7	8	3	.	1	3				8	7	5		
Autophagy	B	N	e	5	A	5	.	3	3	9	3				3	3	4		
protein	—	7	s	.	—	9	4	3	7	0	3	1	2	1	0	0	0	6	2

Atg27,	0	1	C	1	7	0	3												0
putative	1		A	G	.		3												
OS=Neosart	4		D	1	3		3												
orya	8		A	5	3		3												
fumigata	0		F	2	3		3												
(strain	0		U	5															
CEA10 / CBS			B	0															
144.89 /			P																
FGSC				0															
A1163)				0															
GN=AFUB_0				0															
14800 PE=4				0															
SV=1				1															
				4															
				5															
				1															
				5															
				0															
tr B0Y912				8															
B0Y912_AS				5															
PFC				.															
Endonuclea			C																
se/exonucle			A																
ase/phosph			D																
atase family			A																
protein			F	A															
OS=Neosart	A		U	F															
orya	F		B	U															
fumigata	U		P	A															
(strain	B		0	_															
CEA10 / CBS			0	1															
144.89 /	0	B	0	G	4	1													1
FGSC	7	0	0	0	0	7	0												1 9
A1163)	9	Y	7	1	.	.	0												0 0
GN=AFUB_0	2	9	7	5	5	9	4	7	6										9 3
79260 PE=4	6	1	n	2	4	8	.	3	8	2									1 4
SV=1	0	2	o	7	0	9	3	9	8	8	1	0	2	1	0	0	0	0	0 6 0 0 0
tr B0XNLO	A		5	A						8									
B0XNLO_AS	F		0	F						4									
PFC	U		8	U						1									
Extracellular	B		5	A						0									
cell wall	_	B	.	_				4	4										
glucanase	0	0	C	1				0	.										8 8 8
Crf1/allerge	1	X	A	G	0	2	.	6											7 2 2
n Asp F9	5	N	y	D	1	.	2	3	2	6									1 5 5
OS=Neosart	5	L	e	A	6	7	.	9	6	6									7 4 9
orya	3	0	s	F	1	7	5	5	9	7	1	1	1	1	0	0	0	0	3 3 8 0 0 0

fumigata	0	U	9																
(strain		B	0																
CEA10 / CBS		P																	
144.89 /		0																	
FGSC		0																	
A1163)		0																	
GN=AFUB_0		0																	
15530 PE=4		1																	
SV=1		5																	
		2																	
		2																	
		5																	
		0																	
		8																	
		5																	
tr BOY842		.																	
BOY842_AS		C																	
PFC Patched		A																	
sphingolipid		D																	
transporter		A																	
(Ncr1),		F	A																
putative		U	F																
OS=Neosart	A	B	U																
orya	F	P	A																
fumigata	U	0	—																
(strain	B	0	6																
CEA10 / CBS	—	0	G																
144.89 /	0	B	0																
FGSC	7	0	0	0	0		1	8					7	7	9				
A1163)	6	Y	7	9	.		1	4	1				0	7	6				
GN=AFUB_0	0	8	y	4	9	6	4	2	0	3			0	7	4				
76030 PE=4	3	4	e	0	8	5	.	7	.	9			3	0	3				
SV=1	0	2	s	7	0	7	6	3	4	3	2	1	2	3	0	0	0	0	9
																			3
																			7
																			0
tr BOY088			5																0
BOY088_AS			0																0
PFC			8																0
Palmitoyl-			5	A															0
protein	A		.	F															8
thioesterase	F		C	U															0
OS=Neosart	U		A	A															7
orya	B		D	—															4
fumigata	—		A	3						8									4
(strain	0	B	F	G					3	0									4
CEA10 / CBS	3	0	U	1	0				7	4									7
144.89 /	5	Y	B	4	.				.	4									4
FGSC	1	0	P	0	5	5	3	4	6										4
A1163)	5	8	n	0	6	9	.	3	9	.									2
GN=AFUB_0	0	8	o	0	0	2	4	3	5	5	1	2	1	0	0	0	0	0	9
																			4
																			0
																			0
																			0
																			0
																			0
																			0
																			0


```

7
5
0
8
5
.
tr|B0Y6J5|B      C
OY6J5_ASPF      A
C alpha-1,2-     D
Mannosidas      A
e                F A
OS=Neosart      A   U F
orya            F   B U       5
fumigata        U   P A       7
(strain         B   0 _       3
CEA10 / CBS    _   0 4       2
144.89 /       0 B  0 G       7 8
FGSC            6 0  0 1 0     3 .       4 7 5
A1163)         7 Y  6 0 .     . 3       7 4 0
GN=AFUB_0     1 6 y  5 0 6 1 6 6 3     2 1 6
67180 PE=3    8 J e  3 7 4 . 4 0 3     2 2 4
SV=1          0 5 s  7 0 4 6 1 6 3 1 1 1 1 0 0 0 2 0 3 0 0 0

```

<p>Proteins with signal peptide which are quantified in 115146 and not in Ku80 at 30d and 2h of extraction and also at 10d and 2h of extraction (Ranked by average iBAQ intensity in 115146 at 30d 2h)</p>	<p>Statistics (rank from the highest mean of iBAQ values to the lowest to rank from the most abundance)</p>	<p>Number of identified peptides specific to each protein in each sample</p>	<p>iBAQ (intensity-based absolute quantification) values in each sample</p>
--	---	--	---

		nt prot ein to the least)																		
			P																	
			r																	
			o																	
			t																	
							U													
							n													
							n													
							n													
							i													
							i													
							q													
							u													
							e													
							.													
							.													
							p													
							p													
							e													
							p													
							p													
							t													
							i													
							d													
							e													
							e													
							s.													
							s.													
							s.													
							3													
							3													
							0													
							d													
							d													
							2													
							2													
							h													
							h													
							h													
							-													
							-													
							1													
							1													
							1													
							5													
							5													
							4													
							6													
							0													
							0													
							0													
							K													
							K													
							K													
							U													
							U													
							U													
							4													
							4													
							6													
							0													
							0													
							0													
							-													
							-													
							1													
							2													
							3													
							1													
							2													
							3													

	I P i f p r e d i c t e d)				
--	--	--	--	--	--

		5																			
		0																			
		8																			
		5																			
		.																			
tr BOYBK6		C																			
BOYBK6_AS		A																			
PFC		D																			
Uncharacter		A																			
ized protein		F	A																		
OS=Neosart	A	U	F																		
orya	F	B	U																		
fumigata	U	P	A																		
(strain	B	0	_		1												1	1	1		
CEA10 / CBS	_	0	7		7												7	7	8		
144.89 /	0	B	0	G		2	5										5	0	0		
FGSC	8	0	0	0	0		1	7									9	9	5		
A1163)	6	Y	8	0	.	2	.	9									1	4	2		
GN=AFUB_0	9	B	y	4	3	7	9	1	3	0							0	0	0		
86950 PE=4	5	K	e	5	7	8	.	9	8	0							0	0	0		
SV=1	0	6	s	1	0	2	3	8	4	0	4	4	4	4	0	0	0	0	0	0	0
tr BOY1U6	A	5	A		7																
BOY1U6_AS	F	0	F		9																
PFC	U	8	U		3																
Agmatinase,	B	5	A		2												6	8	8		
putative	_	B	.	_		4	3										8	4	5		
OS=Neosart	0	0	C	5	0		2	3									3	5	0		
orya	6	Y	A	G	.	2	.	3									5	8	3		
fumigata	0	1	y	D	1	6	8	3	1	.							5	2	3		
(strain	8	U	e	A	3	7	.	9	7	3							0	0	0		
CEA10 / CBS	9	6	s	F	1	8	4	1	9	3	6	6	6	6	0	0	0	0	0	0	0

144.89 /	0	U	8																	
FGSC		B	0																	
A1163)		P																		
GN=AFUB_0		0																		
60890 PE=3		0																		
SV=1		0																		
		0																		
		5																		
		9																		
		4																		
		8																		
		5																		
		0																		
		8																		
		5																		
tr B0Y1D4		.																		
B0Y1D4_AS		C																		
PFC		A																		
Disulfide		D																		
isomerase		A																		
(TigA),		A																		
putative		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U								1									
fumigata	U	P	A								4									
(strain	B	0	-								.									
CEA10 / CBS	-	0	5					5	3			5	6	5						
144.89 /	0	B	0	G			4	9	3			8	4	6						
FGSC	5	0	0	1	0		0	6	3			2	4	2						
A1163)	9	Y	5	2	.	5	.	5	3			6	9	0						
GN=AFUB_0	8	1	y	8	2	8	4	3	2	5	3			8	2	5				
59890 PE=3	9	D	e	5	6	6	.	6	5	0	3	1	1	1				0	0	0
SV=1	0	4	s	7	0	7	6	8	3	0	3	4	5	4	0	0	0	0	0	0
sp B0Y766				5																
P20D1_ASP				0																
FC Probable				8																
carboxypep				5	A															
tidase	A	.	F																	
AFUB_0727	F	C	U							3	5									
30	U	A	A							4	.									
OS=Neosart	B	D	-							6	6									
orya	-	A	6							6	6			2	3	4				
fumigata	0	B	F	G				4	9	6				6	5	1				
(strain	7	0	U	0	0			6	6	6				7	3	9				
CEA10 / CBS	2	Y	B	6	.	2	.	6	6					3	4	2				
144.89 /	7	7	y	P	8	5	5	4	3	.	6			2	9	8				
FGSC	3	6	e	0	0	8	.	4	7	6	6			0	0	0				
A1163)	0	6	s	0	0	4	9	0	7	7	7	5	6	6	0	0	0	0	0	0

GN=AFUB_0			0																							
72730 PE=3			0																							
SV=1			7																							
			0																							
			8																							
			1																							
			5																							
			0																							
tr BOY110			8																							
BOY110_AS			5																							
PFC			.																							
Secretory pathway protein			C																							
Ssp120, putative			A																							
OS=Neosart orya	A	F	U	F																						
fumigata (strain)	F		B	U						3																
CEA10 / CBS 144.89 / FGSC A1163)	U		P	A						.																
GN=AFUB_0 59210 PE=4	B		0	_						6																
SV=1	_		0	5		2	6			2	3	3														
	0	B	0	G		2	8	6			5	1	0													
	5	0	0	1	0	3	9	6			2	3	2													
	9	Y	5	1	.	3	6				2	3	4													
	2	1	y	7	6	6	2	2	1	6		0	2	1												
	1	1	e	9	4	4	3	0	7	0	6		0	0	0											
	0	0	s	4	0	4	6	3	7	0	7	4	4	3	0	0	0	0	0	0	0	0	0	0	0	0
			5																							
tr BOY4S2			0																							
BOY4S2_AS			8																							
PFC			5																							
Endosomal cargo receptor (Erv25), putative			.																							
OS=Neosart orya			C																							
fumigata (strain)			A																							
CEA10 / CBS 144.89 / FGSC A1163)			D																							
GN=AFUB_0 70080 PE=3	A	F	F																							
SV=1	F		U	U						2	4															
	U		B	A						3	.															
	B		P	_						8	3															
	_		0	4		1	3				2	2	2													
	0	B	0	G		3	5	3			3	0	7													
	7	0	0	1	0	0	6	3			3	7	3													
	0	Y	0	3	.	3	.	6	3		5	8	0													
	0	4	y	6	1	9	3	2	0	.	3	1	8	8												
	8	S	e	8	9	2	.	6	3	6	3	0	0	0												
	0	2	s	2	0	7	8	6	9	7	3	5	3	5	0	0	0	0	0	0	0	0	0	0	0	0

			8														
			5														
			0														
			8														
tr B0YCQ5			5														
B0YCQ5_AS			.														
PFC SUN			C														
domain			A														
protein			D														
(Uth1),			A														
putative			F A														
OS=Neosart	A		U F														
orya	F		B U									5					
fumigata	U		P A									.					
(strain	B		0 _									3					
CEA10 / CBS	_		0 7									2 3					
144.89 /	0 B		0 G									4 1 3					
FGSC	9 0		0 0 0									3 1 3					
A1163)	1 Y		8 5 . 2									. 0 3					
GN=AFUB_0	0 C y		8 4 4 3 4 5 2 3														
91030 PE=4	3 Q e		5 5 9 . 1 0 0 3														
SV=1	0 5 s		0 0 6 2 4 4 0 3 6 4 6 0 0 0 0 0 0 0 0 0 0														
			5														
			0														
			8														
tr B0Y842			5														
B0Y842_AS			.														
PFC Patched			C														
sphingolipid			A														
transporter			D														
(Ncr1),			A														
putative			F A														
OS=Neosart	A		U F														
orya	F		B U														2
fumigata	U		P A									.					
(strain	B		0 _									3					
CEA10 / CBS	_		0 6									1 3					
144.89 /	0 B		0 G									4 3					
FGSC	7 0		0 0 0									1 8 3					
A1163)	6 Y		7 9 .									1 4 2 3					
GN=AFUB_0	0 8 y		4 9 6 4 2 0 5 3														
76030 PE=4	3 4 e		0 8 5 . 7 . 1 3														
SV=1	0 2 s		7 0 7 6 3 4 1 3 3 2 2 0 0 0 0 0 3 0 0 0 0														
tr B0XXV2	A B		5 A 0 3 1 1														4 1 1
B0XXV2_AS	F 0 y		0 F . 7 1 7 4														8 9 9
PFC DNA	U X e		8 U 6 . 5 . 5														7 1 6
damage	B X s		5 A 9 4 5 2 4 3 1 4 4 0 0 0 9 0 6 0 0 0														

response	_	V	.	_	4		2	9			2	5	3
protein	0	2	C	3			1	0			0	0	0
(Dap1),	3		A	G				6			0	0	
putative	8		D	1				.					
OS=Neosart	6		A	0				6					
orya	7		F	4				7					
fumigata	0		U	9									
(strain			B	0									
CEA10 / CBS			P										
144.89 /			0										
FGSC			0										
A1163)			0										
GN=AFUB_0			0										
38670 PE=3			3										
SV=1			7										
			8										
			6										
			5										
			0										
tr BOXNLO			8										
BOXNLO_AS			5										
PFC			.										
Extracellular			C										
cell wall			A										
glucanase			D										
Crf1/allerge			A										
n Asp F9			F	A									
OS=Neosart	A		U	F									
orya	F		B	U									
fumigata	U		P	A									
(strain	B		0	_									
CEA10 / CBS	_		0	1				1			1	1	1
144.89 /	0	B	0	G			4	2			3	0	2
FGSC	1	0	0	1			0	1			2	7	4
A1163)	5	X	1	6	0	2	.	2			0	0	6
GN=AFUB_0	5	N	y	5	1	.	2	3	2	4	3	7	2
15530 PE=4	3	L	e	2	9	7	.	9	6	0	0	0	0
SV=1	0	0	s	2	0	7	5	5	9	0	4	4	4
											4	4	0
											0	0	0
											0	0	0
											0	0	0
											0	0	0
											0	0	0
											0	0	0
tr BOXUX8	A		5	A				2					
BOXUX8_AS	F		0	F				.					
PFC	U		8	U				1	3		2		1
Translocon-	B	B	5	A				2	1	3	0	2	2
associated	_	0	.	_	0			8	6	3	1	8	0
protein,	0	X	C	2	.	4		.	9	3	5	6	7
alpha	3	U	y	A	G	8	2	2	1	6	0	7	2
subunit,	1	X	e	D	1	4	.	6	5	6	0	8	0
putative	8	8	s	A	6	1	1	1	8	0	3	3	1
											3	1	3
											0	0	0
											0	0	0
											0	0	0
											0	0	0
											0	0	0
											0	0	0

GN=AFUB_0		0																		
43810 PE=4		0																		
SV=1		0																		
		4																		
		2																		
		9																		
		9																		
		5																		
		0																		
		8																		
		5																		
		.																		
tr B0XQR8		C																		
B0XQR8_AS		A																		
PFC		D																		
Endopolyph		A																		
osphatase		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U					2												
fumigata	U	P	A					.												
(strain	B	0	_					6												
CEA10 / CBS	_	0	1					6									1			
144.89 /	0	B	0	G			7	8	6							5	0	9		
FGSC	1	0	0	1	0			6	4	6						2	5	6		
A1163)	0	X	1	1	.			.	6	6						5	3	1		
GN=AFUB_0	9	Q	y	0	4	7	3	6	2	9	6					7	6	4		
10920 PE=4	2	R	e	7	9	8	.	6	0	3	6					2	0	7		
SV=1	0	8	s	2	0	2	4	8	6	0	7	2	3	3	0	0	0	0	0	0
		5																		
		0																		
		8																		
		5																		
tr B0XUQ5		5																		
B0XUQ5_AS		.																		
PFC		C																		
Allergen		A																		
Asp F4		D	A																	
OS=Neosart	A	A	F																	
orya	F	F	U					7	2											
fumigata	U	U	A					8	.											
(strain	B	B	_					4	6											
CEA10 / CBS	_	P	2					4	6											
144.89 /	0	B	0	G				3	4	6						3	2	7		
FGSC	2	0	0	0	0			4	3	6						2	6	6		
A1163)	0	X	0	3	.	2		.	.	6						6	1	5		
GN=AFUB_0	9	U	y	0	8	5	5	3	0	3	6					0	4	9		
20900 PE=4	0	Q	e	2	3	0	.	2	9	3	6					1	0	2		
SV=1	0	5	s	0	0	1	2	2	3	3	7	1	4	3	0	0	0	0	0	0

			3																	
			9																	
			5																	
			0																	
			8																	
tr BOYCL1			5																	
BOYCL1_ASP			.																	
FC			C																	
Glyoxylase			A																	
family			D																	
protein,			A																	
putative			F	A																
OS=Neosart	A		U	F																
orya	F		B	U			7	2												
fumigata	U		P	A			4	.												
(strain	B		0	_			6	6												
CEA10 / CBS	_		0	7			2	6												
144.89 /	0	B	0	G			2	3	6					5	9	7				
FGSC	9	0	0	0	0		5	6	6					4	2	7				
A1163)	0	Y	8	5	.	1	.	.	6					6	0	1				
GN=AFUB_0	5	C	y	8	0	5	4	2	2	6	6			0	7	9				
90580 PE=4	8	L	e	0	1	6	.	2	7	6	6			7	4	0				
SV=1	0	1	s	6	5	1	7	5	3	7	7	2	3	3	0	0	0	0	0	0
			5																	
			0																	
			8																	
			5																	
			.																	
tr BOYA75			C																	
BOYA75_AS			A																	
PFC			D																	
Uncharacter			A																	
ized protein			F	A																
OS=Neosart	A		U	F																
orya	F		B	U			6	1												
fumigata	U		P	A			8	.												
(strain	B		0	_			9	6												
CEA10 / CBS	_		0	8			5	6												
144.89 /	0	B	0	G			4	9	6					7	7	4				
FGSC	8	0	0	0	0		5	6	6					9	8	8				
A1163)	3	Y	8	3	.		.	.	6					4	7	6				
GN=AFUB_0	6	A	y	1	9	5	9	4	0	6	6			3	6	8				
83670 PE=4	7	7	e	4	0	6	.	1	9	6	6			7	0	2				
SV=1	0	5	s	4	5	5	6	5	7	7	7	2	2	1	0	0	0	0	0	0
tr BOXTX2	A	B	y	5	A	0	6	2	2	6				3	8	7				
BOXTX2_AS	F	0	e	0	F	.	.	3	6	7				9	3	8				
PFC Cortical	U	X	s	8	U	8	7	9	.	0	1	1	1	1	0	0	0	7	3	1

patch	B T	5 A 2	6 7	9 2 0
protein	_ X	. _ 1	0 2	1 1 5
SUR7,	0 2	C 2	9 3	0 0 0
putative	1	A G	.	
OS=Neosart	9	D 0	3	
orya	4	A 2	3	
fumigata	0	F 3	3	
(strain	0	U 1		
CEA10 / CBS		B 0		
144.89 /		P		
FGSC		0		
A1163)		0		
GN=AFUB_0		0		
19400 PE=4		0		
SV=1		1		
		8		
		9		
		3		
		5		
		0		
		8		
		5		
tr B0XQ77		.		
B0XQ77_AS		C		
PFC Dynein		A		
light chain		D		
(Tctex1),		A		
putative		F A		
OS=Neosart	A	U F		
orya	F	B U	6	
fumigata	U	P A	3	
(strain	B	0 _	9	
CEA10 / CBS	_	0 1	1	
144.89 /	0 B	0 G	1 8	6 5 6
FGSC	0 0	0 0 0	5 6	5 9 6
A1163)	9 X	0 9 . 2	. .	9 7 0
GN=AFUB_0	0 Q y	8 5 5 1 1 3 6		4 6 5
09000 PE=4	0 7 e	8 5 2 . 4 2 6		0 2 4
SV=1	0 7 s	1 0 6 7 3 5 7	2 2 2 2 0 0 0 0 0 0 0 0 0 0 0	
tr B0XQR6	A	5 A		
B0XQR6_AS	F	0 F		
PFC	U B	8 U	2 5	8 3
Endosomal	B 0	5 A 0	5 9	7 0
cargo	_ X	. _ . 2	. 0	2 9
receptor	0 Q y	C 1 7 7 2 0 8		5 2
(P24),	1 R e	A G 8 . 1 5 8		4 3
putative	0 6 s	D 1 3 1 8 6 5	2 3 1 2 0 0 0 0 0 0 0 0 0 0 0	

OS=Neosart	9	A	1																			
orya	0	F	4																			
fumigata	0	U	7																			
(strain		B	0																			
CEA10 / CBS		P																				
144.89 /			0																			
FGSC			0																			
A1163)			0																			
GN=AFUB_0			0																			
10900 PE=3			1																			
SV=1			0																			
			7																			
			0																			
			5																			
			0																			
			8																			
tr B0YDZ9			5																			
B0YDZ9_AS			.																			
PFC Ser/Thr			C																			
protein			A																			
phosphatas			D																			
e family			A																			
protein			F A																			
OS=Neosart	A		U F																			
orya	F		B U		5	4																
fumigata	U		P A		6	.																
(strain	B		0 _		5	6																
CEA10 / CBS	_		0 6		2	6																
144.89 /	0 B		0 G		7	6	6					4	6	6								
FGSC	9 0		0 0 0		5	6	6					7	0	2								
A1163)	7 Y		9 0 . 2		.	.	6					0	4	0								
GN=AFUB_0	3 D y		4 1 7 3		6	5	6	6				9	3	5								
97340 PE=4	4 Z e		6 6 3 .		6	4	6	6				5	4	1								
SV=1	0 9 s		4 0 2 9		6	7	7	7	5	4	5	0	0	0	0	0	0	0	0	0	0	0
tr B0XXQ4			5																			
B0XXQ4_AS			0 A																			
PFC	A		8 F																			
Glutaminas	F		5 U			5																
e, putative	U		. A			4																
OS=Neosart	B		C _			8																
orya	_		A 3			3																
fumigata	0 B		D G		9	4						6	4	5								
(strain	3 0		A 1 0		2	3						5	2	7								
CEA10 / CBS	8 X		F 0 . 1		.	.						1	2	1								
144.89 /	2 X y		U 9 6 7		8	7	3					6	0	2								
FGSC	2 Q e		B 1 5 .		3	5	3					7	7	9								
A1163)	0 4 s		P 0 1 9		4	7	3	6	7	5	6	0	0	0	0	0	0	0	0	0	0	0

GN=AFUB_0 0
 38220 PE=4 0
 SV=1 0
 0
 3
 7
 4
 4

 sp|BOXRS8|
 ECM14_ASP
 FC Putative
 metallocarb
 oxypeptidas
 e ecm14
 OS=Neosart 5 2
 orya 2 .
 fumigata 8 3
 (strain 5 3
 CEA10 / CBS B 6 0 3 1 6 7
 144.89 / e 0 0 6 3 3 6 8 3
 FGSC c X # # . . 3 6 5 3
 A1163) m R y N N 9 6 5 0 3 3 2 8 4
 GN=ecm14 1 S e / / 5 . 8 6 3 3 2 2 7
 PE=3 SV=1 4 8 s A A 1 7 6 8 3 3 3 2 2 0 0 0 0 0 0 0 0 0
 tr|BOY449|
 BOY449_AS 5
 PFC 0
 Dolichyl- 8
 diphosphool 5
 igosaccharid .
 e--protein C
 glycosyltran A
 sferase D
 subunit A
 WBP1 F A
 OS=Neosart A U F
 orya F B U 2
 fumigata U P A .
 (strain B 0 _ 3
 CEA10 / CBS _ 0 5 3
 144.89 / 0 B 0 G 5 3 3 3 2 3
 FGSC 5 0 0 0 1 3 3 3 9 6
 A1163) 6 Y 5 8 0 1 . 1 3 1 4 8
 GN=AFUB_0 5 4 y 5 9 . 6 4 6 6 3 5 5 9
 56510 PE=3 1 4 e 3 7 9 . 6 4 8 3 5 1 8
 SV=1 0 9 s 3 0 4 9 2 2 0 3 3 2 2 0 0 0 0 0 0 0 0 0
 tr|BOY269| A B y 5 A 0 1 3 3 3 2 3 2 2 0 0 0 3 2 3 0 0 0

1
 5
 0
 8
 5
 .
 tr|BOYAM7 C
 |BOYAM7_A A
 SPFC Class V D
 chitinase A
 ChiB1 F A
 OS=Neosart A U F
 orya F B U 0
 fumigata U P A .
 (strain B 0 _ 6
 CEA10 / CBS _ 0 8 6
 144.89 / 0 B 0 G 4 2 6 1 3
 FGSC 8 0 0 0 0 7 4 6 5 3
 A1163) 5 Y 8 1 . . 5 6 7 3
 GN=AFUB_0 2 A y 2 4 7 4 6 5 6 9 0
 85200 PE=3 0 M e 9 1 6 3 2 0 6 3 7
 SV=1 0 7 s 6 0 4 9 3 2 0 7 1 1 0 0 0 0 0 0 0 0 0 0 0 0
 5
 0
 8
 5
 .
 tr|BOY335| C
 BOY335_AS A
 PFC D
 Uncharacter A
 ized protein F A
 OS=Neosart A U F
 orya F B U 2 2
 fumigata U P A 2 .
 (strain B 0 _ 9 3
 CEA10 / CBS _ 0 5 4 3
 144.89 / 0 B 0 G 2 4 3 1 2 2
 FGSC 5 0 0 0 0 6 3 3 7 7 4
 A1163) 2 Y 5 4 . 1 . . 3 1 4 2
 GN=AFUB_0 8 3 y 1 2 8 6 2 1 3 3 3 5 4
 52800 PE=4 0 3 e 6 8 3 . 4 1 3 3 2 9 2
 SV=1 0 5 s 9 0 2 7 5 5 3 3 2 3 2 0 0 0 0 0 0 0 0 0 0
 tr|BOXVA9| A B 5 A 0 1 3 1 1 1 1
 BOXVA9_AS F 0 y 0 F . 7 3 6 6 7 8 4
 PFC Lectin U X e 8 U 6 . 2 . 7 6 2 4
 family B V s 5 A 9 4 8 0 9 2 2 2 2 0 0 0 8 0 8 0 0 0

integral	-	A	.	-	1			7	3									6	9	5
membrane	0	9	C	2					3									0	0	0
protein,	3		A	G					.											
putative	2		D	1					3											
OS=Neosart	4		A	6					3											
orya	7		F	8					3											
fumigata	0		U	0																
(strain			B	0																
CEA10 / CBS			P																	
144.89 /			0																	
FGSC			0																	
A1163)			0																	
GN=AFUB_0			0																	
32470 PE=4			3																	
SV=1			1																	
			8																	
			3																	
			5																	
			0																	
			8																	
			5																	
tr B0XY72			.																	
B0XY72_AS			C																	
PFC Cell			A																	
wall			D																	
glucanase,			A																	
putative			F	A																
OS=Neosart	A		U	F																
orya	F		B	U																
fumigata	U		P	A																
(strain	B		0	-																
CEA10 / CBS	-		0	3																
144.89 /	0	B	0	G				1									1		1	
FGSC	3	0	0	0	0			4	0								0	9	2	
A1163)	9	X	3	9	.			0	7								0	8	3	
GN=AFUB_0	8	Y	y	9	2	9	6	3	.	5							6	0	8	
39870 PE=4	7	7	e	0	5	3	.	6	5	3							8	3	8	
SV=1	0	2	s	6	0	3	3	3	9	3	1	1	1	1	0	0	0	0	9	0
tr B0Y6J5 B	A		5	A																
OY6J5_ASPF	F		0	F															9	
C alpha-1,2-	U		8	U															3	
Mannosidas	B	B	5	A															7	8
e	-	0	.	-	0														4	3
OS=Neosart	0	Y	C	4	.														8	1
orya	6	6	y	A	G	6	1	6	6	3									9	6
fumigata	7	J	e	D	1	4	.	4	0	3									3	1
(strain	1	5	s	A	0	4	6	1	6	3	1	1	1	1	0	0	0	0	9	0

CEA10 / CBS	8	F	0	3
144.89 /	0	U	7	
FGSC		B	0	
A1163)		P		
GN=AFUB_0		0		
67180 PE=3		0		
SV=1		0		
		0		
		6		
		5		
		3		
		7		

<p>Proteins with signal peptide which are quantified in 115146 and not in Ku80 at 10d and 2h of extraction and also at 30d and 2h of extraction</p> <p>(Ranked by average iBAQ intensity in 115146 at 10d 2h)</p>	<p>Statistics (rank from the highest mean of iBAQ values to the lowest to rank from the most abundant protein to the least)</p>	<p>Number of identified peptides specific to each protein in each sample</p>	<p>iBAQ (intensity-based absolute quantification) values</p>
---	---	--	--

Fasta.headers			Prot			With			Presentation							
1	1	5	1	1	4	6	1	1	5	1	1	4	6	1	1	5
2	2	9	2	2	9	3	2	2	9	3	2	2	9	3	2	2
3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
4	4	3	4	4	3	4	4	3	4	4	3	4	4	3	4	4
5	5	3	5	5	3	5	5	3	5	5	3	5	5	3	5	5
6	6	3	6	6	3	6	6	3	6	6	3	6	6	3	6	6
7	7	3	7	7	3	7	7	3	7	7	3	7	7	3	7	7
8	8	3	8	8	3	8	8	3	8	8	3	8	8	3	8	8
9	9	3	9	9	3	9	9	3	9	9	3	9	9	3	9	9
10	10	3	10	10	3	10	10	3	10	10	3	10	10	3	10	10
11	11	3	11	11	3	11	11	3	11	11	3	11	11	3	11	11
12	12	3	12	12	3	12	12	3	12	12	3	12	12	3	12	12
13	13	3	13	13	3	13	13	3	13	13	3	13	13	3	13	13
14	14	3	14	14	3	14	14	3	14	14	3	14	14	3	14	14
15	15	3	15	15	3	15	15	3	15	15	3	15	15	3	15	15
16	16	3	16	16	3	16	16	3	16	16	3	16	16	3	16	16
17	17	3	17	17	3	17	17	3	17	17	3	17	17	3	17	17
18	18	3	18	18	3	18	18	3	18	18	3	18	18	3	18	18
19	19	3	19	19	3	19	19	3	19	19	3	19	19	3	19	19
20	20	3	20	20	3	20	20	3	20	20	3	20	20	3	20	20
21	21	3	21	21	3	21	21	3	21	21	3	21	21	3	21	21
22	22	3	22	22	3	22	22	3	22	22	3	22	22	3	22	22
23	23	3	23	23	3	23	23	3	23	23	3	23	23	3	23	23
24	24	3	24	24	3	24	24	3	24	24	3	24	24	3	24	24
25	25	3	25	25	3	25	25	3	25	25	3	25	25	3	25	25
26	26	3	26	26	3	26	26	3	26	26	3	26	26	3	26	26
27	27	3	27	27	3	27	27	3	27	27	3	27	27	3	27	27
28	28	3	28	28	3	28	28	3	28	28	3	28	28	3	28	28
29	29	3	29	29	3	29	29	3	29	29	3	29	29	3	29	29
30	30	3	30	30	3	30	30	3	30	30	3	30	30	3	30	30
31	31	3	31	31	3	31	31	3	31	31	3	31	31	3	31	31
32	32	3	32	32	3	32	32	3	32	32	3	32	32	3	32	32
33	33	3	33	33	3	33	33	3	33	33	3	33	33	3	33	33
34	34	3	34	34	3	34	34	3	34	34	3	34	34	3	34	34
35	35	3	35	35	3	35	35	3	35	35	3	35	35	3	35	35
36	36	3	36	36	3	36	36	3	36	36	3	36	36	3	36	36
37	37	3	37	37	3	37	37	3	37	37	3	37	37	3	37	37
38	38	3	38	38	3	38	38	3	38	38	3	38	38	3	38	38
39	39	3	39	39	3	39	39	3	39	39	3	39	39	3	39	39
40	40	3	40	40	3	40	40	3	40	40	3	40	40	3	40	40
41	41	3	41	41	3	41	41	3	41	41	3	41	41	3	41	41
42	42	3	42	42	3	42	42	3	42	42	3	42	42	3	42	42
43	43	3	43	43	3	43	43	3	43	43	3	43	43	3	43	43
44	44	3	44	44	3	44	44	3	44	44	3	44	44	3	44	44
45	45	3	45	45	3	45	45	3	45	45	3	45	45	3	45	45
46	46	3	46	46	3	46	46	3	46	46	3	46	46	3	46	46
47	47	3	47	47	3	47	47	3	47	47	3	47	47	3	47	47
48	48	3	48	48	3	48	48	3	48	48	3	48	48	3	48	48
49	49	3	49	49	3	49	49	3	49	49	3	49	49	3	49	49
50	50	3	50	50	3	50	50	3	50	50	3	50	50	3	50	50

		I P i f p r e d i c t e d)					
--	--	--	--	--	--	--	--

tr BOY1D4		5																	
BOY1D4_AS		0																	
PFC		8																	
Disulfide		5																	
isomerase		.																	
(TigA),		C																	
putative		A																	
OS=Neosart	A	A	F	A															
orya	F	U	F																
fumigata	U	B	U																
(strain	B	P	A																
CEA10 / CBS		0	–																
144.89 /	0	0	4																
FGSC	5	0	8																
A1163)	9	0	0																
GN=AFUB_0	8	1	5																
59890 PE=3	9	D	8																
SV=1	0	e	4																
tr BOXUX8	A	s	7																
BOXUX8_AS	F	0	0																
PFC	U	7	6																
Translocon-	B	0	8																
associated	–	1	3																
protein,	0	0	0																
alpha	3	0	0																
subunit,	1	X	2																
putative	8	U	2																
OS=Neosart	0	y	4																
	0	e	0																
	8	s	7																
	F	1	1																
	1	1	1																
	1	8	0																
	4	4	4																
	2	6	0																
	0	0	0																
	0	0	0																
	0	0	0																
	0	0	0																

orya	0	U	2															
fumigata		B	0															
(strain		P																
CEA10 / CBS		0																
144.89 /		0																
FGSC		0																
A1163)		0																
GN=AFUB_0		3																
31800 PE=4		1																
SV=1		1																
		7																
		5																
		0																
tr BOY4S2		8																
BOY4S2_ASP		5																
FC		.																
Endosomal		C																
cargo		A																
receptor		D																
(Erv25),		A																
putative		F	A															
OS=Neosart	A	U	F															
orya	F	B	U		3	5												
fumigata	U	P	A		2	.												
(strain	B	0	_		9	3												
CEA10 / CBS	_	0	4		6	3				3	3	3						
144.89 /	0	B	0	G	3	2	3			1	5	1						
FGSC	7	0	0	1	0	0	6	3		7	9	2						
A1163)	0	Y	6	3	.	3	.	6	3	3	1	4						
GN=AFUB_0	0	4	y	8	1	9	3	2	0	.	3							
70080 PE=3	8	S	e	2	9	2	.	6	3	6	3							
SV=1	0	2	s	8	0	7	8	6	9	7	3	6	6	4	0	0	0	0
tr BOXXV2		5																
BOXXV2_AS		0																
PFC DNA		8																
damage		5	A															
response	A	.	F															
protein	F	C	U		3	4												
(Dap1),	U	A	A		0	.												
putative	B	D	_		2	6												
OS=Neosart	_	A	3		5	6					2	2	3					
orya	0	B	F	G	1	7	6				9	9	1					
fumigata	3	0	U	1	0	7	3	6			5	6	6					
(strain	8	X	B	0	.	3	.	3	6		5	0	1					
CEA10 / CBS	6	X	y	P	4	6	7	1	2	.	6							
144.89 /	7	V	e	0	9	9	.	5	2	3	6							
FGSC	0	2	s	0	0	4	4	5	1	3	7	5	5	4	0	0	0	0

4
 5
 0
 8
 sp|B0Y766| 5
 P20D1_ASP .
 FC Probable C
 carboxypept A
 idase D
 AFUB_0727 A
 30 F A
 OS=Neosart A U F
 orya F B U 1 3
 fumigata U P A 8 .
 (strain B 0 _ 9 6
 CEA10 / CBS _ 0 6 4 6 2 1 1
 144.89 / 0 B 0 G 4 4 6 3 9 3
 FGSC 7 0 0 0 0 6 3 6 3 8 6
 A1163) 2 Y 7 6 . 2 . 3 6 5 2 5
 GN=AFUB_0 7 7 y 0 8 5 5 4 3 . 6 1 5 7
 72730 PE=3 3 6 e 8 0 8 . 4 7 3 6 0 0 0
 SV=1 0 6 s 1 0 4 9 0 7 3 7 4 4 3 0 0 0 0 0 0 0 0 0 0

5
 0
 8
 5
 .
 tr|B0Y1U6| C
 B0Y1U6_AS A
 PFC D
 Agmatinase, A
 putative F A
 OS=Neosart A U F
 orya F B U 1 3
 fumigata U P A 6 .
 (strain B 0 _ 3 3
 CEA10 / CBS _ 0 5 1 3 1 1 1
 144.89 / 0 B 0 G 4 6 3 4 5 8
 FGSC 6 0 0 1 0 2 6 3 5 7 6
 A1163) 0 Y 5 3 . 2 . 6 3 9 4 0
 GN=AFUB_0 8 1 y 9 1 6 8 3 1 . 3 2 9 9
 60890 PE=3 9 U e 4 8 7 . 9 7 6 3 0 0 0
 SV=1 0 6 s 8 0 8 4 1 9 7 3 3 4 3 0 0 0 0 0 0 0 0 0
 tr|B0XQR6| A B 5 A 0 2 2 1 4 1 1 1
 B0XQR6_AS F 0 y 0 F . 7 2 5 5 . 4 7 4
 PFC U X e 8 U 7 . 1 . 9 6 9 9 9
 Endosomal B Q s 5 A 8 1 8 0 5 6 5 5 4 0 0 0 4 7 3 0 0 0

cargo	—	R	.	—	3	5	1	6													6	6	2
receptor	0	6	C	1		6	3	6													0	0	0
(P24),	1		A	G		3	6														0	0	0
putative	0		D	1		.	6																
OS=Neosart	9		A	1		3	6																
orya	0		F	4		3	7																
fumigata	0		U	7																			
(strain			B	0																			
CEA10 / CBS			P																				
144.89 /			0																				
FGSC			0																				
A1163)			0																				
GN=AFUB_0			0																				
10900 PE=3			1																				
SV=1			0																				
			7																				
			0																				
			5																				
			0																				
			8																				
			5																				
tr B0YDZ9			.																				
B0YDZ9_AS			C																				
PFC Ser/Thr			A																				
protein			A																				
phosphatas			D																				
e family			A																				
protein			F	A																			
OS=Neosart	A		U	F																			
orya	F		B	U				1															
fumigata	U		P	A				5															
(strain	B		0	—				6															
CEA10 / CBS	—		0	6				0													1	1	1
144.89 /	0	B	0	G				7	8												4	5	6
FGSC	9	0	0	0	0			5	3												6	8	2
A1163)	7	Y	9	0	.	2	.	3													8	5	9
GN=AFUB_0	3	D	y	4	1	7	3	6	5	.											0	4	1
97340 PE=4	4	Z	e	6	6	3	.	6	4	3	1	1	1								0	0	0
SV=1	0	9	s	4	0	2	9	6	7	3	0	1	0	9	0	0	0	0	0	0	0	0	0
tr B0YCQ5	A		5	A				1	5														
B0YCQ5_AS	F		0	F				4	.														
PFC SUN	U		8	U				7	6												1	1	
domain	B	B	5	A				4	4	6											6	9	7
protein	—	0	.	—	0			3	4	6											6	7	8
(Uth1),	0	Y	C	7	.	2	.	6	6												2	9	1
putative	9	C	y	A	G	4	3	4	5	6	6										1	8	5
OS=Neosart	1	Q	e	D	0	9	.	1	0	.	6										0	0	0
orya	0	5	s	A	5	6	2	4	4	6	6	5	5	7	0	0	0	0	0	0	0	0	0

fumigata	3	F	4		7	7												
(strain	0	U	5															
CEA10 / CBS		B	0															
144.89 /		P																
FGSC		0																
A1163)		0																
GN=AFUB_0		0																
91030 PE=4		0																
SV=1		8																
		8																
		5																
		0																
tr B0Y449																		
B0Y449_AS		5																
PFC		0																
Dolichyl-		8																
diphosphool		5																
igosaccharid		.																
e--protein		C																
glycosyltran		A																
sferase		D																
subunit		A																
WBP1		F	A															
OS=Neosart	A	U	F															
orya	F	B	U					4										
fumigata	U	P	A					.										
(strain	B	0	_					3										
CEA10 / CBS	_	0	5					3								1		
144.89 /	0	B	0	G				5	9	3					9	1	8	
FGSC	5	0	0	0				1	9	3					9	5	3	
A1163)	6	Y	5	8	0	1		.	4	3					5	6	0	
GN=AFUB_0	5	4	y	5	9	.	6	4	6	1	3				4	8	1	
56510 PE=3	1	4	e	3	7	9	.	6	4	3	3				5	0	4	
SV=1	0	9	s	3	0	4	9	2	2	0	3	5	5	3	0	0	0	0
tr B0XQ77																		
B0XQ77_AS		0	A															
PFC Dynein	A	8	F															
light chain	F	5	U					8										
(Tctex1),	U	.	A					6										
putative	B	C	_					5										
OS=Neosart	_	A	1					8										
orya	0	B	D	G				1	8						6	0	9	
fumigata	0	0	A	0	0			5	6						2	0	7	
(strain	9	X	F	9	.	2		.	.						0	3	3	
CEA10 / CBS	0	Q	y	U	5	5	1	1	3	6					5	7	4	
144.89 /	0	7	e	B	5	2	.	4	2	6					3	0	3	
FGSC	0	7	s	P	0	6	7	3	5	7	2	2	2	2	0	0	0	0

A1163)		0																	
GN=AFUB_0		0																	
09000 PE=4		0																	
SV=1		0																	
		0																	
		8																	
		8																	
		1																	
		5																	
		0																	
		8																	
		5																	
		.																	
tr BOYBK6		C																	
BOYBK6_AS		A																	
PFC		D																	
Uncharacter		A																	
ized protein		F A																	
OS=Neosart	A	U F																	
orya	F	B U					7												
fumigata	U	P A					9												
(strain	B	0 _					0												
CEA10 / CBS	_	0 7					7												
144.89 /	0 B	0 G					2 7								7 7 8				
FGSC	8 0	0 0 0					1 6								9 7 0				
A1163)	6 Y	8 0 . 2					. .								8 0 3				
GN=AFUB_0	9 B y	4 3 7 9					1 3 6								6 7 0				
86950 PE=4	5 K e	5 7 8 . 9 8 6													0 1 2				
SV=1	0 6 s	1 0 2 3 8 4 7					1 1 1 1 0 0 0 0 0 0 0 0												
		5																	
		0																	
tr BOYA75		8																	
BOYA75_AS		5																	
PFC		.																	
Uncharacter		C																	
ized protein		A A																	
OS=Neosart	A	D F																	
orya	F	A U					7												
fumigata	U	F A					6												
(strain	B	U _					1												
CEA10 / CBS	_	B 8					6												
144.89 /	0 B	P G					4 0								6 7 9				
FGSC	8 0	0 0 0					5 3								0 4 4				
A1163)	3 Y	0 3 .					. .								0 3 0				
GN=AFUB_0	6 A y	0 9 5 9					4 0 3								6 6 4				
83670 PE=4	7 7 e	0 0 6 . 1 9 3													7 5 9				
SV=1	0 5 s	8 5 5 6 5 7 3					3 3 3 3 0 0 0 0 0 0 0 0												

									1																				
									4																				
									4																				
									5																				
									0																				
									8																				
									5																				
tr BOYCL1									.																				
BOYCL1_ASP									C																				
FC									A																				
Glyoxylase									A																				
family									D																				
protein,									A																				
putative									F	A																			
OS=Neosart	A								U	F																			
orya	F								B	U	6	1																	
fumigata	U								P	A	7	.																	
(strain	B								0	—	6	6																	
CEA10 / CBS	—								0	7	8	6																	
144.89 /	0	B							0	G	2	0	6					8	7	3									
FGSC	9	0							0	0	0	5	3	6				8	4	9									
A1163)	0	Y							8	5	.	1	.	.	6			6	4	9									
GN=AFUB_0	5	C	y						8	0	5	4	2	2	3	6			0	4	9								
90580 PE=4	8	L	e						0	1	6	.	2	7	3	6			2	2	7								
SV=1	0	1	s						6	5	1	7	5	3	3	7	2	2	1	0	0	0	0	0	0	0	0	0	
									5																				
									0																				
									8																				
tr BOXVA9									5																				
BOXVA9_AS									.																				
PFC Lectin									C																				
family									A																				
integral									D																				
membrane									A																				
protein,									F	A																			
putative									U	F																			
OS=Neosart	A								B	U	5	3																	
orya	F								P	A	4	.																	
fumigata	U								0	—	0	6																	
(strain	B								0	2	2	6																	
CEA10 / CBS	—								0	G	7	6																	
144.89 /	0	B							0	1	0	3	3	6						6	5	4							
FGSC	3	0							3	6	.	1	6	.	6					1	8	1							
A1163)	2	X							1	8	6	7	3	.	3	6				6	8	5							
GN=AFUB_0	4	V	y						8	0	9	.	2	0	3	6				4	6	6							
32470 PE=4	7	A	e						3	0	1	4	8	7	3	7	4	4	3	0	0	0	0	0	0	0	0	0	0
SV=1	0	9	s						5	A	0	1	1	1	5	1				6	3	5							
tr BOXRG2	A	B	y						0	F	.	2	7	8	3	.	2	1	1	0	0	0	8	6	6	0	0	0	
BOXRG2_AS	F	0	e																										

PFC IgE	U X s	8 U 5 . 9 . 8 3		4 5 6
binding	B R	5 A 1 3	5 7 3	0 8 3
protein,	_ G	. _ 1	1 2 3	5 1 1
putative	0 2	C 1	4 3 3	0 0 0
OS=Neosart	0	A G	. 3	
orya	2	D 0	3 3	
fumigata	3	A 1	3 3	
(strain	6	F 9	3 3	
CEA10 / CBS	0	U 8		
144.89 /		B 0		
FGSC		P		
A1163)		0		
GN=AFUB_0		0		
02360 PE=4		0		
SV=1		0		
		0		
		2		
		3		
		1		
		5		
		0		
		8		
		5		
tr BOXTX2		.		
BOXTX2_AS		C		
PFC Cortical		A		
patch		D		
protein		A		
SUR7,		F A		
putative		U F		
OS=Neosart	A	B U		
orya	F	P A		
fumigata	U	0 _		
(strain	B	0 2		
CEA10 / CBS	_	0 G	2 5	4 3 7
144.89 /	0 B	0 0 0	6 1	8 3 2
FGSC	1 0	1 2 .	. 6	6 9 3
A1163)	9 X	8 3 8 6 2 6 6		7 2 9
GN=AFUB_0	4 T y	9 1 2 . 3 0 3		1 4 4
19400 PE=4	0 X e	3 0 1 7 9 9 0	1 1 1 1 0 0 0	0 0 0 0 0 0 0
SV=1	0 2 s	5 A		
tr BOXY72	A	0 F	5	4 6 4
BOXY72_AS	F B	8 U 0	4 1	6 0 6
PFC Cell	U 0	5 A .	0 4	8 7 7
wall	B X	. _ 9 6 3 . 6		5 5 9
glucanase,	_ Y y	C 3 3 . 6 5 5		0 0 5
putative	0 7 e	A G 3 3 3 9 0	2 2 2 2 0 0 0	0 0 0 0 0 0 0
OS=Neosart	3 2 s			

orya	9	D	0																	
fumigata	8	A	9																	
(strain	7	F	2																	
CEA10 / CBS	0	U	5																	
144.89 /		B	0																	
FGSC		P																		
A1163)		0																		
GN=AFUB_0		0																		
39870 PE=4		0																		
SV=1		0																		
		3																		
		9																		
		0																		
		6																		
		5																		
		0																		
		8																		
		5																		
		.																		
tr BOXZB5		C																		
BOXZB5_AS		A																		
PFC Ser/Thr		D																		
protein		A																		
phosphatas		A																		
e family		F	A																	
OS=Neosart	A	U	F																	
orya	F	B	U				1													
fumigata	U	P	A				.													
(strain	B	0					3													
CEA10 / CBS		0	3				3													
144.89 /	0	B	0	G			7	3	3					5	2	3				
FGSC	4	0	0	0	0		1	7	3					5	4	3				
A1163)	3	X	4	4	.	1	.	7	3					6	5	1				
GN=AFUB_0	8	Z	y	2	1	5	4	6	2	7	3			6	3	0				
43810 PE=4	1	B	e	9	6	1	.	2	6	0	3			9	6	5				
SV=1	0	5	s	9	0	4	4	6	2	0	3	2	1	1	0	0	0	0	0	0
sp BOXRS8																				
ECM14_ASP																				
FC Putative																				0
metallocarb																				.
oxypeptidas																				6
e ecm14																				6
OS=Neosart	B						6	3	6											2
orya	e	0				0	6	0	6											6
fumigata	c	X				#	#	.	.	8	6									2
(strain	m	R	y			N	N	9	6	5	0	8	6							8
CEA10 / CBS	1	S	e			/	/	5	.	8	6	1	6							4
144.89 /	4	8	s			A	A	1	7	6	8	0	7	1	1	0	0	0	0	0

FGSC
 A1163)
 GN=ecm14
 PE=3 SV=1

		5																			
		0																			
		8																			
		5																			
		.																			
tr BOXR8		C																			
BOXR8_AS		A																			
PFC		D																			
Endopolyph		A																			
osphatase		F A																			
OS=Neosart	A	U F																			
orya	F	B U					2														
fumigata	U	P A					9														
(strain	B	0 _					5														
CEA10 / CBS	_	0 1					4														
144.89 /	0 B	0 G					7 7										2 2 4				
FGSC	1 0	0 1 0					6 3										5 0 2				
A1163)	0 X	1 1 .					. .										6 8 0				
GN=AFUB_0	9 Q y	0 4 7 3					6 2 3										9 6 7				
10920 PE=4	2 R e	7 9 8 .					6 0 3										9 9 4				
SV=1	0 8 s	2 0 2 4					8 6 3 2 2 2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0														

		5																				
		0																				
		8																				
		5																				
		.																				
tr BOY335		C																				
BOY335_AS		A																				
PFC		D																				
Uncharacter		A																				
ized protein		F A																				
OS=Neosart	A	U F																				
orya	F	B U					2 2															
fumigata	U	P A					9 .															
(strain	B	0 _					3 3															
CEA10 / CBS	_	0 5					0 3															
144.89 /	0 B	0 G					2 2 3										2 2 3					
FGSC	5 0	0 0 0					6 6 3										6 4 6					
A1163)	2 Y	5 4 . 1					. . 3										3 6 9					
GN=AFUB_0	8 3 y	1 2 8 6 2 1 6 3															5 5 0					
52800 PE=4	0 3 e	6 8 3 . 4 1 6 3															0 0 8					
SV=1	0 5 s	9 0 2 7 5 5 7 3					2 2 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0															
tr BOY269	A B y	5 A 0 1 3 3 1 1 1 1 2 0 0 0 1 1 1 0 0 0 0 0																				

BOY269_AS	F 0 e	0 F . 2 5 8 6 .							4 8 7
PFC	U Y s	8 U 8 . 9 . 9 3							8 6 5
NlpC/P60- like cell-wall peptidase, putative	B 2	5 A 0 3	7 8 3						2 0 3
	_ 6	. _ 9	6 9 3						9 4 6
	0 9	C 5	3 6 3						0 0 0
	6	A G	. 3						
OS=Neosart orya	1	D 1	6 3						
	4	A 3	6 3						
fumigata (strain	7	F 7	7 3						
	0	U 3							
CEA10 / CBS 144.89 / FGSC A1163)		B 0							
		P							
		0							
		0							
GN=AFUB_0 61470 PE=4 SV=1;tr BOY 9E0 BOY9E0 _ASPFC Uncharacter ized protein OS=Neosart orya fumigata (strain CEA10 / CBS 144.89 /		0							
		6							
		0							
		0							
		4							
		5							
		0							
		8							
		5							
tr BOYAM7 BOYAM7_AS PFC Class V chitinase ChiB1 OS=Neosart orya fumigata (strain CEA10 / CBS 144.89 / FGSC A1163) GN=AFUB_0 85200 PE=3 SV=1		.							
		C							
		A							
		D							
		A A							
	A	F F							
	F	U U			1				
	U	B A			5				
	B	P _			9				
	_	0 8			9				
	0 B	0 G	4 8						1 2 1
	8 0	0 0 0	7 6						4 0 3
	5 Y	0 1 .	. .						0 3 6
	2 A y	8 4 7	4 6 6						2 0 6
	0 M e	2 1 6	3 2 6						5 4 7
	0 7 s	9 0 4 9	3 2 7	1 1 1 1 0 0 0 0 0 0 0 0					

				6																
				5																
				0																
				8																
				5																
				.																
				C																
tr BOXUQ5				A																
BOXUQ5_AS				D																
PFC				A																
Allergen				F	A															
Asp F4				U	F															
OS=Neosart	A			B	U			1	1											
orya	F			P	A			3	.											
fumigata	U			0	_			8	3											
(strain	B			0	2			8	3											
CEA10 / CBS	_			0	G			3	3	3					2					
144.89 /	0	B		0	0	0		4	6	3					1	9	9			
FGSC	2	0		2	3	.	2	.	.	3					9	8	8			
A1163)	0	X		0	8	5	5	3	0	6	3				7	0	7			
GN=AFUB_0	9	U	y	3	3	0	.	2	9	6	3				6	5	0			
20900 PE=4	0	Q	e	9	0	1	2	2	3	7	3	2	1	1	0	0	0	0	0	0
SV=1	0	5	s	5																
				0																
				8																
				5																
tr BOXN71				.																
BOXN71_AS				C																
PFC				A																
Autophagy				D																
protein				A																
Atg27,				F	A															
putative				U	F															
OS=Neosart	A			B	U			1	1											
orya	F			P	A			2	.											
fumigata	U			0	_			1	3											
(strain	B			0	1			9	3											
CEA10 / CBS	_			0	G			3	0	3					2					
144.89 /	0	B		0	0	0		7	0	3					6	0	8			
FGSC	1	0		1	5	.		.	.	3					9	7	8			
A1163)	4	X		4	2	7	8	3	3	3	3				8	3	5			
GN=AFUB_0	8	N	y	5	5	5	.	3	7	3	3				3	2	4			
14800 PE=4	0	7	e	1	0	9	4	3	7	3	3	1	2	1	0	0	0	6	0	5
SV=1	0	1	s	1	0	9	4	3	7	3	3	1	2	1	0	0	0	6	0	5
tr BOXNLO	A	B		5	A	0	2	4	8						8	8	8			
BOXNLO_AS	F	0	y	0	F	.	2	3	0	4					7	2	2			
PFC	U	X	e	8	U	7	.	9	.	1					1	5	5			
Extracellular	B	N	s	5	A	7	5	5	2	0	1	1	1	1	0	0	0	7	4	9

cell wall	-	L	.	-	6	4												3	3	8					
glucanase	0	0	C	1	9	.																			
Crf1/allerge	1		A	G		6																			
n Asp F9	5		D	1		6																			
OS=Neosart	5		A	6		6																			
orya	3		F	1		7																			
fumigata	0		U	9																					
(strain			B	0																					
CEA10 / CBS			P																						
144.89 /			0																						
FGSC			0																						
A1163)			0																						
GN=AFUB_0			0																						
15530 PE=4			1																						
SV=1			5																						
			2																						
			2																						
			5																						
			0																						
			8																						
			5																						
tr BOY842			.																						
BOY842_AS			C																						
PFC Patched			A																						
sphingolipid			D																						
transporter			A																						
(Ncr1),			A																						
putative			F	A																					
OS=Neosart	A		U	F																					
orya	F		B	U																					
fumigata	U		P	A																					
(strain	B		0	-																					
CEA10 / CBS	-		0	6																					
144.89 /	0	B	0	G																					
FGSC	7	0	0	0	0	1	8											7	7	9					
A1163)	6	Y	7	9	.	1	4	1										0	7	6					
GN=AFUB_0	0	8	y	4	9	6	4	2	0	3								0	7	4					
76030 PE=4	3	4	e	0	8	5	.	7	.	9								3	0	3					
SV=1	0	2	s	7	0	7	6	3	4	3	2	1	2	3	0	0	0	0	9	3	7	0	0	0	
tr BOY6J5 B	A		5	A																					
OY6J5_ASPF	F		0	F																					
C alpha-1,2-	U		8	U																					
Mannosidas	B	B	5	A																					
e	-	0	.	-	0																				
OS=Neosart	0	Y	C	4	.																				
orya	6	6	y	A	G	6	1	6	6	3															
fumigata	7	J	e	D	1	4	.	4	0	3															
(strain	1	5	s	A	0	4	6	1	6	3	1	1	1	1	0	0	0	0	2	0	3	0	0	0	

CEA10 / CBS	8	F	0	3
144.89 /	0	U	7	
FGSC		B	0	
A1163)		P		
GN=AFUB_0		0		
67180 PE=3		0		
SV=1		0		
		0		
		6		
		5		
		3		
		7		

ACCEPTED MANUSCRIPT




Figure 1. Structure (A), internal dynamics (B) and hydrophobicity (C) of monomeric RodA (best energy conformer). **A:** Strands are shown in magenta, helices in cyan and disulfide bridges in marine blue. The inter cysteine regions on the front (grey) and back (light grey) as well as the N-terminal region are indicated. Cysteines are numbered by their order in the sequence throughout this paper. **B:** Backbone amide ^1H - ^{15}N heteronuclear nOe values are shown on the cartoon representation of the structure: highly flexible ($n\text{Oe} < 0.45$), flexible ($0.45 \leq n\text{Oe} \leq 0.65$), rigid ($n\text{Oe} > 0.65$) and missing-value backbone residues are shown in red, orange, blue and grey, respectively. The structure is shown in a different orientation on the right. **C:** The hydropathy profile calculated with the Eisenberg hydrophobicity scale ([http:// web.expasy.org/protscale](http://web.expasy.org/protscale)) is color coded on the surface and cartoon representations of the structure: blue= hydrophilic ($\Phi \leq -0.3$), orange ($-0.3 < \Phi < 0.3$), red=hydrophobic ($\Phi \geq 0.3$). The N-terminal disordered region (residues 18-38) is not displayed in **B** and **C**.




Figure 2. The effect of mutation(s) on the kinetics of RodA rodlet self-assembly assessed by Thioflavin T fluorescence at 50 °C. Amyloid assembly profiles for (A) RodA WT; single point mutants (B) RodA I114G, (C) RodA I115G, (D) RodA I145G and (E) RodA I146G; (F) double point mutant RodA I115G-I146G and (G) C7C8NC2 chimera. (H) Lag time and (I) time taken to reach half maximum ThT fluorescence. (See the material and methods section for calculation of lag time and time to reach half maximum).




Figure 3. Assembly of RodA WT and mutant proteins on highly oriented pyrolytic graphite. Atomic force micrographs showing the surface morphology of rodlets formed by (A) RodA WT, (B) RodA I114G, (C) RodA I115G, (D) RodA I145G and (E) RodA I146G after incubation at room temperature on graphite surfaces. Surface layers containing rodlets formed by (F) RodA WT, (G) double mutant RodA I115G/I146G and (H) C7C8NC2 chimera after incubation at 50 °C for 2 hours. (See the material and methods section for the AFM experimental conditions).




Figure 4. Transmission electron micrographs of fibrillar material formed by (A) PIIGIQDL, a peptide derived from the C4-C5 region of RodA and (B) SLIGL, a peptide with predicted amyloidogenic sequence from the C7-C8 region.

ACCEPTED MANUSCRIPT

ACCEPTED




Figure 5. Cysteine mutations impair the presence of rodlets at the conidial surface. AFM deflection images showing the structure of conidial surface for: (A) the parental, (B) one cysteine (C127S) and (C) four cysteines (C64S/C65S/C133S/C134S) mutants.




Figure 6. Delay in the rodlets appearance at the surface of conidial mutants. AFM deflection images of the rodlets layer for (A-C) the parental, (D-F) I146G and (G-I) I115S/I146G mutants after 3 (A, D, G), 10 (B, E, H) and 30 (C, F, I) days of growth.




Figure 7. The surface of the RodA I115S/I146G double mutant is only partially covered by rodlets. AFM deflections images (at different magnification) representative of the conidium surface of the RodA I115S/I146G double mutant after 30 days of culture with different magnifications.

MANUSCRIPT

ACCEPTED




Figure 8. Recruitment of cells in the lungs of C57/Bl6 mice after inhalation of conidia of the single RodA I115S, double RodA I115S/I146G, $\Delta rodA$ mutant and the parental strain Ku80. Total number of cells were counted in the BALs (counts from > 6 mice per experiment, 2 experiments) 6 hours after conidial inoculation and were expressed as average with SEM. No significant differences were observed between the two mutants and the parental strain, while more cells were significantly recruited in the lungs of mice infected with the $\Delta rodA$ mutant.




Figure 9. Conidia of single and quadruple cystein mutants and the double RodA I115S/I146G mutant activate the human immune response.

A-B: Induction of the dendritic cells (DC) maturation. **C-D:** Production of cytokines by DC. **E-F:** Polarization of T cell response. As other single point mutants did not alter either DC phenotype nor cytokines, they were not tested for T cell response. Note that (i) the cystein mutants also polarizes the Treg response and that (ii) the I146G mutant slightly activates the DC response.




Figure 10. Lack of DC activation by the reduced (A,B) or mutated RodA protein (C,D). A-B: Unfolded rRodA, with or without reduced and blocked Cys residues with N-ethyl maleimide (NEM) or iodoacetamide (IA) do not activate dendritic cells (A) and do not induce cytokine production (B). C-D: RodA extracted from conidia of mutants L145S, I146G, I115G/I146S (C) or lacking the N terminal region (RodA-Nterm) (D), are immunologically inert as parental RodA. LPS is used as a control.




Figure 11. Degradation of rRodA by the major proteases identified in *A. fumigatus*, Alp1, Mep1, Pep1 and Pep2. rRodA (4 µg) was mixed with different proteases (1 µg or less as indicated) in 30 µl of 20 mM Tris-HCl pH 7 for Alp1 and Mep1 or sodium acetate pH 5.5, 4 or 3 for Pep1 and Pep2. Samples were taken after 1.5 h or 2 h of incubation at 37 °C. PMSF, EDTA or slightly acidic pH were used as inhibitors of Alp1, Mep1 and Pep1/2 respectively.